arXiv:0812.5003v1 [math.RA] 30 Dec 2008

Topological $N=2$ superconformal superbialgebras

Lifang Lin[†)], Huanxia Fa[‡)], Jianhua Zhou[*,†)]

[†)]Department of Mathematics, Southeast University, Nanjing 210096, China

[‡)]Department of Mathematics, Changshu Institute of Technology, Changshu 215500, China

Abstract. Lie superbialgebra structures on the centerless topological $N=2$ superconformal algebra $\mathcal{T}$ are considered, all of which are proved to be coboundary triangular.

Key words: Lie super-bialgebras, Yang-Baxter equation, topological $N=2$ conformal algebra.

Mathematics Subject Classification (2000): 17B62, 17B05, 17B37, 17B66.

§1. Introduction

The notion of Lie bialgebras was first introduced by Drinfeld in 1983 in a connection with Hamiltonian mechanics and Possion Lie group. Lie bialgebras arising from non-abelian two dimensional Lie subalgebras of Lie algebras (such as Witt, one-sided Witt, and Virasoro algebras) have been considered in [3, 4]. It is well known that the $N=2$ superconformal algebras were first independently constructed by Kac (see [2]) and by Ademollo et al., which play important roles in both mathematics and physics. The determination of Lie bialgebra structures on a Lie algebra is considered to be the first partial step towards quantizing Lie algebras. Lie superbialgebra structures on the Ramond and twisted $N=2$ superconformal algebra were determined in [5] and [6] respectively. In the present paper, we study Lie superbialgebra structures on the centerless topological $N=2$ superconformal algebra. It is proved that all such Lie superbialgebras are coboundary triangular.

Firstly we shall recall some related concepts. Let $\mathcal{T}=\mathcal{T}^{\bar{0}}\oplus\mathcal{T}^{\bar{1}}$ be a super-vector space over the complex number field $\mathbb{C}$. All elements in the following are assumed to be $\mathbb{Z}_2$-homogeneous. For $x\in\mathcal{T}$, the notation $[x]$ means $x\in\mathcal{T}^{[x]}$. Denote by τ and ξ the *super-twist map* of $\mathcal{L}\otimes\mathcal{L}$ and *super-cyclic map* of $\mathcal{L}\otimes\mathcal{L}\otimes\mathcal{L}$, i.e., $\tau(x_1\otimes x_2)=(-1)^{[x_1][x_2]}x_2\otimes x_1$, $\xi=(1\otimes\tau)\cdot(\tau\otimes 1): x_1\otimes x_2\otimes x_3\mapsto(-1)^{[x_1]([x_2]+[x_3])}x_2\otimes x_3\otimes x_1$ for $x_1,x_2,x_3\in\mathcal{L}$.

A *Lie superalgebra* is a pair $(\mathcal{T},\varphi)$ consisting of a super-vector space $\mathcal{T}$ and a bilinear map $\varphi:\mathcal{T}\otimes\mathcal{T}\to\mathcal{T}$ such that $\varphi(\mathcal{L}^i,\mathcal{L}^j)\subset\mathcal{L}^{i+j}$, $\mathrm{Ker}(1\otimes 1-\tau)\subset\mathrm{Ker}\,\varphi$, $\varphi\cdot(1\otimes\varphi)\cdot(\mathbf{1}+\xi+\xi^2)=0$. A *Lie super-coalgebra* is a pair $(\mathcal{T},\Delta)$ consisting of a super-vector space $\mathcal{T}$ and a linear map $\Delta:\mathcal{T}\to\mathcal{T}\otimes\mathcal{T}$ such that $\Delta(\mathcal{L}^i)\subset\sum\limits_{j+k=i}\mathcal{L}^j\otimes\mathcal{L}^k$, $\mathrm{Im}\,\Delta\subset\mathrm{Im}(1\otimes 1-\tau)$, $(\mathbf{1}+\xi+\xi^2)\cdot(1\otimes\Delta)\cdot\Delta=0$. A *Lie super-bialgebra* is a triple $(\mathcal{T},\varphi,\Delta)$ with $(\mathcal{T},\varphi)$ being a Lie superalgebra, $(\mathcal{T},\Delta)$ a Lie super-coalgebra and $\Delta\varphi(x\otimes y)=x\cdot\Delta y-(-1)^{[x][y]}y\cdot\Delta x$ for $x,y\in\mathcal{T}$. The symbol "$\cdot$"

Supported by NSF grants 10671027 of China

[*]Corresponding author: jhZhou@seu.edu.cn

means the adjoint diagonal action and $[x,y]=\varphi(x\otimes y)$, i.e.,

$$x\cdot(\sum_i a_i\otimes b_i)=\sum_i([x,a_i]\otimes b_i+(-1)^{[x][a_i]}a_i\otimes[x,b_i])\text{ for }x,a_i,b_i\in\mathcal{T}.$$

A *coboundary Lie super-bialgebra* is a quadruple $(\mathcal{T},\varphi,\Delta,r)$, where $(\mathcal{T},\varphi,\Delta)$ is a Lie super-bialgebra and $\Delta=\Delta_r$ is a *coboundary of* r for $r\in\mathrm{Im}(1\otimes 1-\tau)\subset\mathcal{T}\otimes\mathcal{T}$. And Δ_r is defined by $\Delta_r(x)=(-1)^{[r][x]}x\cdot r,\ \forall\ x\in\mathcal{T}$. If $r=\sum_i a_i\otimes b_i$, then $[r]=[a_i]+[b_i]$. A coboundary bialgebra is called *triangular* if it satisfies the *classical Yang-Baxter Equation*:

$$c(r):=[r^{12},r^{13}]+[r^{12},r^{23}]+[r^{13},r^{23}]=0,\tag{1.1}$$

where $r=\sum_i a_i\otimes b_i$, $r^{12}=\sum_i a_i\otimes b_i\otimes 1$, $r^{13}=\sum_i a_i\otimes 1\otimes b_i$ and $r^{23}=\sum_i 1\otimes a_i\otimes b_i$.

Let $V=V^{\bar 0}\oplus V^{\bar 1}$ be a $\mathcal{T}$-module for the superalgebra $\mathcal{T}=\mathcal{T}^{\bar 0}\oplus\mathcal{T}^{\bar 1}$. A $\mathbb{Z}_2$-homogenous linear map $d:\mathcal{T}\to V$ is called a *homogenous derivations of parity* $[d]\in\mathbb{Z}_2$, if $d(\mathcal{L}^i)\subset V^{i+[d]}$ and there exists $[d]$ such that $d([x,y])=(-1)^{[d][x]}x\cdot d(y)-(-1)^{[y]([d]+[x])}y\cdot d(x)$ for $x,y\in\mathcal{T}$. The derivation d is called *even* if $[d]=\bar 0$, *odd* if $[d]=\bar 1$. Denote by $\mathrm{Der}^p(\mathcal{T},V)$ the set of homogenous derivations of parity p. Let $\mathrm{Der}(\mathcal{T},V)=\mathrm{Der}^{\bar 0}(\mathcal{T},V)\oplus\mathrm{Der}^{\bar 1}(\mathcal{T},V)$ be the set of derivations from $\mathcal{T}$ to V and $\mathrm{Inn}(\mathcal{T},V)=\mathrm{Inn}^{\bar 0}(\mathcal{T},V)\oplus\mathrm{Inn}^{\bar 1}(\mathcal{T},V)$ the set of inner derivations, where $\mathrm{Inn}^p(\mathcal{T},V)$ is the set of *inner derivations of parity* p consisting of a_{inn}, $a\in V^p$, defined by $a_{\mathrm{inn}}:x\mapsto(-1)^{[a][x]}x\cdot a$ for $x\in\mathcal{T}$, $[a]=p$.

Denote by $H^1(\mathcal{T},V)$ the *first cohomology group* of $\mathcal{T}$ with coefficients in V. It is known $H^1(\mathcal{T},V)\cong\mathrm{Der}(\mathcal{T},V)/\mathrm{Inn}(\mathcal{T},V)$. Say $r\in\mathcal{T}$ satisfy the *modified Yang-Baxter equation* if

$$x\cdot c(r)=0,\quad\forall\ x\in\mathcal{T}.\tag{1.2}$$

We shall investigate the topological $N=2$ superconformal algebra, $\mathcal{T}=\mathcal{T}^{\bar 0}\oplus\mathcal{T}^{\bar 1}$, generated by the bosonic operators $\mathcal{L}_n$, $\mathcal{H}_n$ and the fermionic operators $\mathcal{G}_n$, $\mathcal{Q}_n$ with the following non-vanishing brackets ($\mathcal{T}^{\bar 0}=\mathrm{Span}_{\mathbb{C}}\{\mathcal{L}_n,\mathcal{H}_n\,|\,n\in\mathbb{Z}\}$ and $\mathcal{T}^{\bar 1}=\mathrm{Span}_{\mathbb{C}}\{\mathcal{G}_n,\mathcal{Q}_n|\,n\in\mathbb{Z}\}$)

$$\begin{aligned}
&[\mathcal{L}_m,\mathcal{Q}_n]=-n\mathcal{Q}_{m+n}, &&\\
&[\mathcal{L}_m,\mathcal{H}_n]=-n\mathcal{H}_{m+n}, &&[\mathcal{H}_m,\mathcal{G}_n]=\mathcal{G}_{m+n},\\
&[\mathcal{L}_m,\mathcal{G}_n]=(m-n)\mathcal{G}_{m+n}, &&[\mathcal{H}_m,\mathcal{Q}_n]=-\mathcal{Q}_{m+n},\\
&[\mathcal{L}_m,\mathcal{L}_n]=(m-n)\mathcal{L}_{m+n}, &&[\mathcal{G}_m,\mathcal{Q}_n]=2\mathcal{L}_{m+n}-2n\mathcal{H}_{m+n}.
\end{aligned}$$

Theorem 1.1. *Every Lie super-bialgebra structure on the the centerless topological $N=2$ superconformal algebra $\mathcal{T}$ is triangular coboundary.*

§3. Proof of main results

The following lemma can be obtained by employing similar techniques as [3, 5].

Lemma 3.1. *(i) The triple* $(\mathcal{T},[\cdot,\cdot],\Delta_r)$ *is a super-bialgebra if and only if* r *satisfies (1.2).*
(ii) If $r\in\mathcal{T}\otimes\mathcal{T}$ *such that* $x\cdot r=0$, $\forall\ x\in\mathcal{T}$, *then* $r=0$.
(iii) An element $r\in\mathrm{Im}(1\otimes 1-\tau)$ *satisfies (1.1) if only if it satisfies (1.2).*

Proposition 3.2. $\mathrm{Der}(\mathcal{T},V)=\mathrm{Inn}(\mathcal{T},V)$, *where* $V=\mathcal{T}\otimes\mathcal{T}$.

Proof. Note that $V=\oplus_{i\in\mathbb{Z}}V_i$ is $\mathbb{Z}$-graded with $V_i=\sum_{j+k=i}\mathcal{T}_j\otimes\mathcal{T}_k$. We say a derivation $d\in\mathrm{Der}(\mathcal{T},V)$ is *homogeneous of degree* $i\in\mathbb{Z}$ if $d(V_j)\subset V_{i+j}$, $\forall\ j\in\mathbb{Z}$ and denote such i by $\deg d$. Set $\mathrm{Der}(\mathcal{T},V)_i=\{d\in\mathrm{Der}(\mathcal{T},V)\,|\,\deg d=i\}$. For $d\in\mathrm{Der}(\mathcal{T},V)$ and any $u\in\mathcal{T}_j$, write $d(u)=\sum_{k\in\mathbb{Z}}v_k\in V$ and set $d_i(v_j)=v_{i+j}$. Then $d_i\in\mathrm{Der}(\mathcal{T},V)_i$ and

$$d=\sum_{i\in\mathbb{Z}}d_i,\quad d_i\in\mathrm{Der}(\mathcal{T},V)_i.\tag{3.1}$$

Such a sum is *summable*, i.e., for any $u\in\mathcal{T}$ and $d(u)=\sum_{i\in\mathbb{Z}}d_i(u)$, there are only finitely many $d_i(u)\neq 0$. All the sums appear in the following are supposed to be summable.

Claim 1. $d_i\in\mathrm{Inn}(\mathcal{T},V)$ *and* $d_0(L_0)=0$, $\forall\ i\in\mathbb{Z}^*$.

For any $i\in\mathbb{Z}^*$, denote $u=-\frac{1}{i}d_i(L_0)\in V_i$. For any $x_j\in\mathcal{T}_j$, applying d_i to $[L_0,x_j]=-jx_j$, using $d_i(x_j)\in V_{i+j}$ and the action of L_0 on V_{i+j} is the scalar $-i-j$, we have

$$-(i+j)d_i(x_j)-(-1)^{[d_j][x_j]}x_j\cdot d_i(L_0)=-jd_i(x_j),\tag{3.2}$$

i.e., $d_i(x_j)=u_{\mathrm{inn}}(x_j)$. Thus $d_i=u_{\mathrm{inn}}$ is inner. Using (3.2) with $i=0$, we obtain $x\cdot d_0(L_0)=0$. Thus by Lemma 3.1, one has $d_0(L_0)=0$.

Claim 2. *By replacing* d_0 *by* d_0-u_{inn} *for some* $u\in V_0$, *one can suppose* $d_0=0$.

For any $n\in\mathbb{Z}^*$, one can write

$$\begin{aligned}d_0(\mathcal{L}_n)=\sum_{i\in\mathbb{Z}}(&a_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{H}_{-i}+a^{\dagger}_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{L}_{-i}+b_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{G}_{-i}+b^{\dagger}_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{L}_{-i}\\&+c_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{Q}_{-i}+c^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{L}_{-i}+d_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{G}_{-i}+d^{\dagger}_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{H}_{-i}\\&+e_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{Q}_{-i}+e^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{H}_{-i}+f_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{Q}_{-i}+f^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{G}_{-i}\\&+\alpha_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{L}_{-i}+\beta_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{H}_{-i}+\gamma_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{G}_{-i}+\mu_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{Q}_{-i}).\end{aligned}$$

Using the following facts, $\mathcal{L}_1\cdot(\mathcal{X}_i\otimes\mathcal{Z}_{-i})=i\mathcal{X}_i\otimes\mathcal{Z}_{1-i}-i\mathcal{X}_{i+1}\otimes\mathcal{Z}_{-i}$ for $\mathcal{X},\mathcal{Z}=\mathcal{H}$ or $\mathcal{Q}$,

$$\begin{aligned}&\mathcal{L}_1\cdot(\mathcal{X}_i\otimes\mathcal{Z}_{-i})=(1-i)\mathcal{X}_{i+1}\otimes\mathcal{Z}_{-i}+i\mathcal{X}_i\otimes\mathcal{Z}_{1-i}\ \text{for}\ \mathcal{X},\mathcal{Z}=\mathcal{L}\ \text{or}\ \mathcal{G},\\&\mathcal{L}_1\cdot(\mathcal{X}_i\otimes\mathcal{Z}_{-i})=(1-i)\mathcal{X}_{i+1}\otimes\mathcal{Z}_{-i}+i\mathcal{X}_i\otimes\mathcal{Z}_{1-i}\ \text{for}\ \mathcal{X}=\mathcal{L}\ \text{or}\ \mathcal{G},\ \mathcal{Z}=\mathcal{H}\ \text{or}\ \mathcal{Q},\\&\mathcal{L}_1\cdot(\mathcal{X}_i\otimes\mathcal{Z}_{-i})=(1+i)\mathcal{X}_i\otimes\mathcal{Z}_{1-i}-i\mathcal{X}_{i+1}\otimes\mathcal{Z}_{-i}\ \text{for}\ \mathcal{X}=\mathcal{H}\ \text{or}\ \mathcal{Q},\ \mathcal{Z}=\mathcal{L}\ \text{or}\ \mathcal{G},\end{aligned}$$

by subtracting a proper linear combination of the above inner derivations, one can rewrite $d_0(\mathcal{L}_1)-\mu_{1,-1}\mathcal{Q}_0\otimes\mathcal{Q}_1-\mu_{1,0}\mathcal{Q}_1\otimes\mathcal{Q}_0$ as

$$\begin{aligned}&a_{1,-1}\mathcal{L}_0\otimes\mathcal{H}_1+a_{1,1}\mathcal{L}_2\otimes\mathcal{H}_{-1}+a^{\dagger}_{1,-2}\mathcal{H}_{-1}\otimes\mathcal{L}_2+a^{\dagger}_{1,0}\mathcal{H}_1\otimes\mathcal{L}_0+b_{1,-2}\mathcal{L}_{-1}\otimes\mathcal{G}_2+b_{1,1}\mathcal{L}_2\otimes\mathcal{G}_{-1}\\&+b^{\dagger}_{1,-2}\mathcal{G}_{-1}\otimes\mathcal{L}_2+b^{\dagger}_{1,1}\mathcal{G}_2\otimes\mathcal{L}_{-1}+c_{1,-1}\mathcal{L}_0\otimes\mathcal{Q}_1+c_{1,1}\mathcal{L}_2\otimes\mathcal{Q}_{-1}+c^{\dagger}_{1,-2}\mathcal{Q}_{-1}\otimes\mathcal{L}_2+c^{\dagger}_{1,0}\mathcal{Q}_1\otimes\mathcal{L}_0\\&+d_{1,-2}\mathcal{H}_{-1}\otimes\mathcal{G}_2+d_{1,0}\mathcal{H}_1\otimes\mathcal{G}_0+d^{\dagger}_{1,-1}\mathcal{G}_0\otimes\mathcal{H}_1+d^{\dagger}_{1,1}\mathcal{G}_2\otimes\mathcal{H}_{-1}+e_{1,-1}\mathcal{H}_0\otimes\mathcal{Q}_1+e_{1,0}\mathcal{H}_1\otimes\mathcal{Q}_0\\&+e^{\dagger}_{1,-1}\mathcal{Q}_0\otimes\mathcal{H}_1+e^{\dagger}_{1,0}\mathcal{Q}_1\otimes\mathcal{H}_0+f_{1,-1}\mathcal{G}_0\otimes\mathcal{Q}_1+f_{1,1}\mathcal{G}_2\otimes\mathcal{Q}_{-1}+f^{\dagger}_{1,-2}\mathcal{Q}_{-1}\otimes\mathcal{G}_2+f^{\dagger}_{1,0}\mathcal{Q}_1\otimes\mathcal{G}_0\\&+\alpha_{1,-2}\mathcal{L}_{-1}\otimes\mathcal{L}_2+\alpha_{1,1}\mathcal{L}_2\otimes\mathcal{L}_{-1}+j_{1,-1}\mathcal{H}_0\otimes\mathcal{H}_1+j_{1,0}\mathcal{H}_1\otimes\mathcal{H}_0+k_{1,-2}\mathcal{G}_{-1}\otimes\mathcal{G}_2+k_{1,1}\mathcal{G}_2\otimes\mathcal{G}_{-1}.\end{aligned}$$

Applying d_0 to $[\mathcal{L}_1,\mathcal{L}_{-1}]=2\mathcal{L}_0$, we have

$$\mathcal{L}_1\cdot d_0(\mathcal{L}_{-1})=\mathcal{L}_{-1}\cdot d_0(\mathcal{L}_1).\tag{3.3}$$

Comparing the coefficients of $\mathcal{L}_i\otimes\mathcal{H}_{-i}$ in (3.3), we obtain $(i-2)a_{-1,i}=(i+1)a_{-1,i+1}$ for $i\neq\pm1,0,2$, which together with the fact that the set $\{i\mid a_{-1,i}\neq0\}$ is of finite rank, implies

$$a_{-1,i}=0\ \text{ for }\ i\neq0,1,2\ \text{ and }\ a_{-1,1}=-2a_{-1,0}=-2a_{-1,2},\ a_{1,-1}=a_{1,1}=0.$$

Applying the same techniques to $\mathcal{L}_i\otimes\mathcal{Q}_{-i}$, $\mathcal{G}_i\otimes\mathcal{H}_{-i}$ and $\mathcal{G}_i\otimes\mathcal{Q}_{-i}$, one has

$$x_{-1,i}=0,\ i\neq0,1,2,\ x_{-1,1}=-2x_{-1,0}=-2x_{-1,2},\ x_{1,-1}=x_{1,1}=0\ \text{ for }\ x=c,f,d^{\dagger}.$$

Comparing the coefficients of $\mathcal{H}_i\otimes\mathcal{L}_{-i}$ in (3.3), one has $(i-1)a^{\dagger}_{-1,i}=(i+2)a^{\dagger}_{-1,i+1}$ for $i\neq-2,0,\pm1$, which implies $a^{\dagger}_{-1,i}=0$ for $i\neq0,\pm1$, $2a^{\dagger}_{-1,-1}=2a^{\dagger}_{-1,1}=-a^{\dagger}_{-1,0}$ and $a^{\dagger}_{1,0}=a^{\dagger}_{1,-2}=0$. Comparing the coefficients of $\mathcal{Q}_i\otimes\mathcal{L}_{-i}$, $\mathcal{H}_i\otimes\mathcal{G}_{-i}$ and $\mathcal{Q}_i\otimes\mathcal{G}_{-i}$ in (3.3), one has

$$y_{-1,i}=0,\ i\neq0,\pm1\ \text{ and }\ 2y_{-1,-1}=2y_{-1,1}=-y_{-1,0},\quad y_{1,0}=y_{1,-2}=0\ \text{ for }\ y=d,c^{\dagger},f^{\dagger}.$$

Comparing the coefficients of $\mathcal{L}_i\otimes\mathcal{G}_{-i}$ and $\mathcal{G}_i\otimes\mathcal{L}_{-i}$, we obtain

$$z_{-1,i}=z_{-1,1}+z_{-1,0}=3z_{-1,-1}+z_{-1,0}+3z_{1,-2}=3z_{-1,2}-z_{-1,0}+3z_{1,1}=0,$$

for $i\neq\pm1,0,2$ and $z=b$ or $z=b^{\dagger}$. Finally, comparing the coefficients of $\mathcal{H}_i\otimes\mathcal{Q}_{-i}$, $\mathcal{Q}_i\otimes\mathcal{H}_{-i}$, $\mathcal{L}_i\otimes\mathcal{L}_{-i}$, $\mathcal{G}_i\otimes\mathcal{G}_{-i}$, $\mathcal{H}_i\otimes\mathcal{H}_{-i}$ and $\mathcal{Q}_i\otimes\mathcal{Q}_{-i}$, we obtain $x_{-1,i}=x_{-1,1}+x_{-1,0}+x_{1,-1}+x_{1,0}=0$,

$$y_{-1,j}=y_{-1,1}+y_{-1,0}=3y_{-1,-1}+y_{-1,0}+3y_{1,-2}=3y_{-1,2}-y_{-1,0}+3y_{1,1}=0,$$

for $i\neq0,1$ and $x=e,\beta,e^{\dagger},\mu$, $j\neq\pm1,0,2$ and $y=\alpha,\gamma$. Then $d_0(\mathcal{L}_1)$ can be rewritten as

$$\begin{aligned}&b_{1,-2}\mathcal{L}_{-1}\otimes\mathcal{G}_2+b_{1,1}\mathcal{L}_2\otimes\mathcal{G}_{-1}+b^{\dagger}_{1,-2}\mathcal{G}_{-1}\otimes\mathcal{L}_2+b^{\dagger}_{1,1}\mathcal{G}_2\otimes\mathcal{L}_{-1}+e_{1,-1}\mathcal{H}_0\otimes\mathcal{Q}_1+e_{1,0}\mathcal{H}_1\otimes\mathcal{Q}_0\\&+e^{\dagger}_{1,-1}\mathcal{Q}_0\otimes\mathcal{H}_1+e^{\dagger}_{1,0}\mathcal{Q}_1\otimes\mathcal{H}_0+\alpha_{1,-2}\mathcal{L}_{-1}\otimes\mathcal{L}_2+\alpha_{1,1}\mathcal{L}_2\otimes\mathcal{L}_{-1}+\beta_{1,-1}\mathcal{H}_0\otimes\mathcal{H}_1\\&+\beta_{1,0}\mathcal{H}_1\otimes\mathcal{H}_0+\gamma_{1,-2}\mathcal{G}_{-1}\otimes\mathcal{G}_2+\gamma_{1,1}\mathcal{G}_2\otimes\mathcal{G}_{-1}+\mu_{1,-1}\mathcal{Q}_0\otimes\mathcal{Q}_1+\mu_{1,0}\mathcal{Q}_1\otimes\mathcal{Q}_0.\end{aligned}$$

For convenience, we introduce the following notations, $u_1 = \mathcal{L}_1 \otimes \mathcal{H}_{-1} - \mathcal{L}_0 \otimes \mathcal{H}_0$, $u_2 = \mathcal{H}_{-1} \otimes \mathcal{L}_1 - \mathcal{H}_0 \otimes \mathcal{L}_0$, $u_3 = \mathcal{L}_1 \otimes \mathcal{Q}_{-1} - \mathcal{L}_0 \otimes \mathcal{Q}_0$, $u_4 = \mathcal{Q}_{-1} \otimes \mathcal{L}_1 - \mathcal{Q}_0 \otimes \mathcal{L}_0$, $u_5 = \mathcal{G}_1 \otimes \mathcal{H}_{-1} - \mathcal{G}_0 \otimes \mathcal{H}_0$, $u_6 = \mathcal{H}_{-1} \otimes \mathcal{G}_1 - \mathcal{H}_0 \otimes \mathcal{G}_0$, $u_7 = \mathcal{G}_1 \otimes \mathcal{Q}_{-1} - \mathcal{G}_0 \otimes \mathcal{Q}_0$, $u_8 = \mathcal{Q}_{-1} \otimes \mathcal{G}_1 - \mathcal{Q}_0 \otimes \mathcal{G}_0$. Observing

$$\begin{aligned}
&\mathcal{L}_{-1} \cdot u_1 = \mathcal{L}_{-1} \otimes \mathcal{H}_0 - 2\mathcal{L}_0 \otimes \mathcal{H}_{-1} + \mathcal{L}_1 \otimes \mathcal{H}_{-2},\ \mathcal{L}_{-1} \cdot u_2 = -2\mathcal{H}_{-1} \otimes \mathcal{L}_0 + \mathcal{H}_{-2} \otimes \mathcal{L}_1 + \mathcal{H}_0 \otimes \mathcal{L}_{-1},\\
&\mathcal{L}_{-1} \cdot u_3 = \mathcal{L}_{-1} \otimes \mathcal{Q}_0 - 2\mathcal{L}_0 \otimes \mathcal{Q}_{-1} + \mathcal{L}_1 \otimes \mathcal{Q}_{-2},\ \mathcal{L}_{-1} \cdot u_4 = -2\mathcal{Q}_{-1} \otimes \mathcal{L}_0 + \mathcal{Q}_{-2} \otimes \mathcal{L}_1 + \mathcal{Q}_0 \otimes \mathcal{L}_{-1},\\
&\mathcal{L}_{-1} \cdot u_5 = \mathcal{G}_{-1} \otimes \mathcal{H}_0 - 2\mathcal{G}_0 \otimes \mathcal{H}_{-1} + \mathcal{G}_1 \otimes \mathcal{H}_{-2},\ \mathcal{L}_{-1} \cdot u_6 = -2\mathcal{H}_{-1} \otimes \mathcal{G}_0 + \mathcal{H}_{-2} \otimes \mathcal{G}_1 + \mathcal{H}_0 \otimes \mathcal{G}_{-1},\\
&\mathcal{L}_{-1} \cdot u_7 = \mathcal{G}_{-1} \otimes \mathcal{Q}_0 - 2\mathcal{G}_0 \otimes \mathcal{Q}_{-1} + \mathcal{G}_1 \otimes \mathcal{Q}_{-2},\ \mathcal{L}_{-1} \cdot u_8 = -2\mathcal{Q}_{-1} \otimes \mathcal{G}_0 + \mathcal{Q}_{-2} \otimes \mathcal{G}_1 + \mathcal{Q}_0 \otimes \mathcal{G}_{-1},
\end{aligned}$$

and $\mathcal{L}_1 \cdot u_i = \mathcal{L}_{-1} \cdot u_i = 0$, $\forall\ i = 1, \cdots, 8$, one can replace d_0 by $d_0 - \vartheta_{\rm inn}$ (where ϑ is a proper linear combination of u_i, $i = 1, \cdots, 8$) and rewrite $d_0(\mathcal{L}_{-1})$ as

$$\begin{aligned}
&b_{-1,0}\mathcal{L}_{-1} \otimes \mathcal{G}_0 - b_{-1,0}\mathcal{L}_0 \otimes \mathcal{G}_{-1} - (b_{-1,0}/3 + b_{1,-2})\mathcal{L}_{-2} \otimes \mathcal{G}_1 + (b_{-1,0}/3 - b_{1,1})\mathcal{L}_1 \otimes \mathcal{G}_{-2}\\
&+\gamma_{-1,0}\mathcal{G}_{-1} \otimes \mathcal{G}_0 - (\gamma_{-1,0}/3 + \gamma_{1,-2})\mathcal{G}_{-2} \otimes \mathcal{G}_1 - \gamma_{-1,0}\mathcal{G}_0 \otimes \mathcal{G}_{-1} + (\gamma_{-1,0}/3 - \gamma_{1,1})\mathcal{G}_1 \otimes \mathcal{G}_{-2}\\
&+b^\dagger_{-1,0}\mathcal{G}_{-1} \otimes \mathcal{L}_0 - b^\dagger_{-1,0}\mathcal{G}_0 \otimes \mathcal{L}_{-1} - (b^\dagger_{-1,0}/3 + b^\dagger_{1,-2})\mathcal{G}_{-2} \otimes \mathcal{L}_1 + (b^\dagger_{-1,0}/3 - b^\dagger_{1,1})\mathcal{G}_1 \otimes \mathcal{L}_{-2}\\
&+\alpha_{-1,0}\mathcal{L}_{-1} \otimes \mathcal{L}_0 - (\alpha_{-1,0}/3 + \alpha_{1,-2})\mathcal{L}_{-2} \otimes \mathcal{L}_1 - \alpha_{-1,0}\mathcal{L}_0 \otimes \mathcal{L}_{-1} + (\alpha_{-1,0}/3 - \alpha_{1,1})\mathcal{L}_1 \otimes \mathcal{L}_{-2}\\
&+e_{-1,0}\mathcal{H}_{-1} \otimes \mathcal{Q}_0 - (e_{-1,0} + e_{1,-1} + e_{1,0})\mathcal{H}_0 \otimes \mathcal{Q}_{-1} + e^\dagger_{-1,0}\mathcal{Q}_{-1} \otimes \mathcal{H}_0 - (e^\dagger_{-1,0} + e^\dagger_{1,-1} + e^\dagger_{1,0})\mathcal{Q}_0 \otimes \mathcal{H}_{-1}\\
&+\beta_{-1,0}\mathcal{H}_{-1} \otimes \mathcal{H}_0 - (\beta_{-1,0} + \beta_{1,-1} + \beta_{1,0})\mathcal{H}_0 \otimes \mathcal{H}_{-1} + \mu_{-1,0}\mathcal{Q}_{-1} \otimes \mathcal{Q}_0 - (\mu_{-1,0} + \mu_{1,-1} + \mu_{1,0})\mathcal{Q}_0 \otimes \mathcal{Q}_{-1}.
\end{aligned}$$

Applying d_0 to $[\mathcal{L}_1, \mathcal{L}_{-2}] = 3\mathcal{L}_{-1}$, one has

$$\mathcal{L}_1 \cdot d_0(\mathcal{L}_{-2}) = \mathcal{L}_{-2} \cdot d_0(\mathcal{L}_1) + 3d_0(\mathcal{L}_{-1}). \tag{3.4}$$

Comparing the coefficients of $\mathcal{L}_{i-1} \otimes \mathcal{H}_{-i}$ in (3.4), one has $(i-3)a_{-2,i} = (i+1)a_{-2,i+1}$ for $i \neq 0, 1, 2$, which together with $\{i \mid \alpha_{-2,i} \neq 0\}$ being finite, implies

$$a_{-2,i} = a_{-2,1} + 3a_{-2,0} = a_{-2,2} - 3a_{-2,0} = a_{-2,3} + a_{-2,0} = 0, \quad \forall\ i \neq 0, 1, 2, 3.$$

Similarly, comparing the coefficients of $\mathcal{L}_{i-1} \otimes \mathcal{Q}_{-i}$, $\mathcal{G}_{i-1} \otimes \mathcal{H}_{-i}$, $\mathcal{G}_{i-1} \otimes \mathcal{Q}_{-i}$, we obtain

$$x_{-2,i} = x_{-2,1} + 3x_{-2,0} = x_{-2,2} - 3x_{-2,0} = x_{-2,3} + x_{-2,0} = 0,\ \forall\ i \neq 0, 1, 2, 3,\ x = c, f, d^\dagger.$$

Comparing the coefficients of $\mathcal{H}_{i-1} \otimes \mathcal{L}_{-i}$ in (3.4), one has $(i-2)a^\dagger_{-2,i} = (i+2)a^\dagger_{-2,i+1}$ for $i \neq \pm 1, 0$, which implies $a^\dagger_{-2,i} = a^\dagger_{-2,1} + a^\dagger_{-2,0} = 3a^\dagger_{-2,-1} + a^\dagger_{-2,0} = 3a^\dagger_{-2,2} - a^\dagger_{-2,0} = 0$, $\forall\ i \neq \pm 1, 0, 2$. Then comparing the coefficients of $\mathcal{Q}_{i-1} \otimes \mathcal{L}_{-i}$, $\mathcal{H}_{i-1} \otimes \mathcal{G}_{-i}$, $\mathcal{Q}_{i-1} \otimes \mathcal{G}_{-i}$, we obtain $y_{-2,i} = y_{-2,1} + y_{-2,0} = 3y_{-2,-1} + y_{-2,0} = 3y_{-2,2} - y_{-2,0} = 0$, $\forall\ i \neq \pm 1, 0, 2$, $y = d, c^\dagger, f^\dagger$. Comparing the coefficients of $\mathcal{L}_{i-1} \otimes \mathcal{G}_{-i}$ and $\mathcal{G}_{i-1} \otimes \mathcal{L}_{-i}$ in (3.4), one has

$$(i+2)z_{-2,i+1} = (i-3)z_{-2,i} \quad \text{for } i \neq \pm 2, \pm 1, 0, 3,\ \ z = b, b^\dagger,$$

which implies $z_{-2,i}=2z_{-2,1}+3z_{-2,0}-3z_{-1,0}=z_{-2,2}-z_{-2,0}+2z_{-1,0}=4z_{-2,-1}+z_{-2,0}+z_{-1,0}=4z_{-2,3}+z_{-2,0}-3z_{-1,0}=z_{1,-2}=z_{1,1}=0$, for all $i\neq\pm1,0,2,3$ and $z=b,b^{\dagger}$. Comparing the coefficients of $\mathcal{H}_{i-1}\otimes\mathcal{Q}_{-i}$ and $\mathcal{Q}_{i-1}\otimes\mathcal{H}_{-i}$ in (3.4), one has $(i-2)e_{-2,i}=(i+1)e_{-2,i+1}$ for $i\neq0,1,\ x=e,e^{\dagger}$, which implies $x_{-2,i}=x_{-2,1}+2x_{-2,0}-3x_{-1,0}+x_{1,0}=x_{-2,2}-x_{-2,0}+3x_{-1,0}+2x_{1,-1}+x_{1,0}=0$ for all $i\neq\pm1,0,2$ and $x=e,e^{\dagger}$. Comparing the coefficients of $\mathcal{L}_{i-1}\otimes\mathcal{L}_{-i}$, $\mathcal{G}_{i-1}\otimes\mathcal{G}_{-i}$, $\mathcal{H}_{i-1}\otimes\mathcal{H}_{-i}$ and $\mathcal{Q}_{i-1}\otimes\mathcal{Q}_{-i}$ in (3.4), we obtain

$$\begin{aligned}
y_{-2,i}&=2y_{-2,1}+3y_{-2,0}-3y_{-1,0}=y_{-2,2}-y_{-2,0}+2y_{-1,0}\\
&=4y_{-2,3}+y_{-2,0}-3y_{-1,0}=4y_{-2,-1}+y_{-2,0}+y_{-1,0}=y_{1,1}=y_{1,-2}=0,\\
z_{-2,j}&=z_{-2,1}+2z_{-2,0}+z_{1,0}-3z_{-1,0}=z_{-2,2}-z_{-2,0}+2z_{1,-1}+3z_{-1,0}+z_{1,0}=0,
\end{aligned}$$

where $i\neq\pm1,0,2,3$, $j\neq0,1,2$, $y=\alpha,\gamma$ and $z=\beta,\mu$. From $d_0([\mathcal{L}_2,\mathcal{L}_{-1}])=3d_0\mathcal{L}_1$, one has

$$\mathcal{L}_{-1}\cdot d_0(\mathcal{L}_2)=\mathcal{L}_2\cdot d_0(\mathcal{L}_{-1})-3d_0(\mathcal{L}_1). \tag{3.5}$$

Comparing the coefficients of $\mathcal{L}_{i+1}\otimes\mathcal{H}_{-i}$ in (3.5), one has $(i+3)a_{2,i}=(i-1)a_{2,i-1}$ for $i\neq\pm1,0,2$, which together with $\{i\mid\alpha_{2,i}\neq0\}$ being finite, implies

$$a_{2,i}=a_{2,-1}+3a_{2,0}=a_{2,-2}-3a_{2,0}=a_{2,-3}+a_{2,0}=0,\ \forall\ i\neq-3,\cdots,0.$$

Similarly, comparing the coefficients of $\mathcal{L}_{i+1}\otimes\mathcal{Q}_{-i}$, $\mathcal{G}_{i+1}\otimes\mathcal{H}_{-i}$, $\mathcal{G}_{i+1}\otimes\mathcal{Q}_{-i}$, we obtain

$$x_{2,i}=x_{2,-1}+3x_{2,0}=x_{2,-2}-3x_{2,0}=x_{2,-3}+x_{2,0}=0,\ \forall\ i\neq-3,\cdots,0,\ x=c,f,d^{\dagger}.$$

Comparing the coefficients of $\mathcal{H}_{i+1}\otimes\mathcal{L}_{-i}$, $\mathcal{Q}_{i+1}\otimes\mathcal{L}_{-i}$, $\mathcal{H}_{i+1}\otimes\mathcal{G}_{-i}$ and $\mathcal{Q}_{i+1}\otimes\mathcal{G}_{-i}$, one has

$$y_{2,i}=3y_{2,1}+y_{2,0}=y_{2,-1}+y_{2,0}=3y_{2,-2}-y_{2,0}=0,\ \forall\ i\neq0,\pm1,-2,\ y=a^{\dagger},c^{\dagger},d,f^{\dagger}.$$

Comparing the coefficients of $\mathcal{L}_{i+1}\otimes\mathcal{G}_{-i}$ and $\mathcal{G}_{i+1}\otimes\mathcal{L}_{-i}$ in (3.5), we obtain

$$z_{2,i}=z_{-1,0}=2z_{2,-1}+3z_{2,0}=4z_{2,1}+z_{2,0}=4z_{2,-3}+z_{2,0}=0,\ z_{2,-2}=z_{2,0},$$

for all $i\neq0,\pm1,-3$ and $z=b,b^{\dagger}$. Comparing the coefficients of $\mathcal{H}_{i+1}\otimes\mathcal{Q}_{-i}$ and $\mathcal{Q}_{i+1}\otimes\mathcal{H}_{-i}$ in (3.5), one has $e_{2,i}=e_{2,-1}+2e_{2,0}+e_{-1,0}-3e_{1,0}=e_{2,-2}-e_{2,0}-e_{-1,0}-2e_{1,-1}+e_{1,0}=0$, for all $i\neq0,-1,-2$ and $x=e,e^{\dagger}$. Comparing the coefficients of $\mathcal{L}_{i+1}\otimes\mathcal{L}_{-i}$ and $\mathcal{G}_{i+1}\otimes\mathcal{G}_{-i}$ in (3.5), one has $y_{2,i}=y_{-1,0}=2y_{2,-1}+3y_{2,0}=y_{2,-2}-y_{2,0}=4y_{2,-3}+y_{2,0}=4y_{2,1}+y_{2,0}=0$, for all $i\neq-3,-2,0,\pm1$ and $y=\alpha,\gamma$. Comparing the coefficients of $\mathcal{H}_{i+1}\otimes\mathcal{H}_{-i}$ and $\mathcal{Q}_{i+1}\otimes\mathcal{Q}_{-i}$ in (3.5), one has $z_{2,i}=z_{2,-1}+2z_{2,0}+z_{-1,0}-3z_{1,0}=z_{2,-2}-z_{2,0}-z_{-1,0}-2z_{1,-1}+z_{1,0}=0$, for all $i\neq-2,-1,0$ and $z=\beta,\mu$. Denote $v_1=2\mathcal{L}_0\otimes\mathcal{G}_0-\mathcal{L}_1\otimes\mathcal{G}_{-1}-\mathcal{L}_{-1}\otimes\mathcal{G}_1$, $v_2=2\mathcal{G}_0\otimes\mathcal{L}_0-$

$\mathcal{G}_1\otimes\mathcal{L}_{-1}-\mathcal{G}_{-1}\otimes\mathcal{L}_1$, $v_3=2\mathcal{L}_0\otimes\mathcal{L}_0-\mathcal{L}_1\otimes\mathcal{L}_{-1}-\mathcal{L}_{-1}\otimes\mathcal{L}_1$, $v_4=2\mathcal{G}_0\otimes\mathcal{G}_0-\mathcal{G}_1\otimes\mathcal{G}_{-1}-\mathcal{G}_{-1}\otimes\mathcal{G}_1$. Observing the facts, $\mathcal{L}_{-2}\cdot v_1=-4\mathcal{L}_{-2}\otimes\mathcal{G}_0+6\mathcal{L}_{-1}\otimes\mathcal{G}_{-1}-4\mathcal{L}_0\otimes\mathcal{G}_{-2}+\mathcal{L}_{-3}\otimes\mathcal{G}_1+\mathcal{L}_1\otimes\mathcal{G}_{-3}$,

$$\begin{aligned}
\mathcal{L}_{-2}\cdot v_2&=-4\mathcal{G}_{-2}\otimes\mathcal{L}_0+6\mathcal{G}_{-1}\otimes\mathcal{L}_{-1}-4\mathcal{G}_0\otimes\mathcal{L}_{-2}+\mathcal{G}_{-3}\otimes\mathcal{L}_1+\mathcal{G}_1\otimes\mathcal{L}_{-3},\\
\mathcal{L}_{-2}\cdot v_3&=-4\mathcal{L}_{-2}\otimes\mathcal{L}_0+6\mathcal{L}_{-1}\otimes\mathcal{L}_{-1}-4\mathcal{L}_0\otimes\mathcal{L}_{-2}+\mathcal{L}_{-3}\otimes\mathcal{L}_1+\mathcal{L}_1\otimes\mathcal{L}_{-3},\\
\mathcal{L}_{-2}\cdot v_4&=-4\mathcal{G}_{-2}\otimes\mathcal{G}_0+6\mathcal{G}_{-1}\otimes\mathcal{G}_{-1}-4\mathcal{G}_0\otimes\mathcal{G}_{-2}+\mathcal{G}_{-3}\otimes\mathcal{G}_1+\mathcal{G}_1\otimes\mathcal{G}_{-3}
\end{aligned}$$

and $\mathcal{L}_1\cdot v_i=\mathcal{L}_{-1}\cdot v_i=0$, $\forall\ i=1,\cdots,4$, one can replace d_0 by $d_0-\varpi_{\rm inn}$ (where ϖ is a proper linear combination of v_i $(i=1,\cdots,4)$) and rewrite $d_0(\mathcal{L}_{-2})$ as

$$\begin{aligned}
&e_{-2,0}\mathcal{H}_{-2}\otimes\mathcal{Q}_0-(2e_{-2,0}-3e_{-1,0}+e_{1,0})\mathcal{H}_{-1}\otimes\mathcal{Q}_{-1}+(e_{-2,0}-3e_{-1,0}-2e_{1,-1}-e_{1,0})\mathcal{H}_0\otimes\mathcal{Q}_{-2}\\
&+e^{\dagger}_{-2,0}\mathcal{Q}_{-2}\otimes\mathcal{H}_0-(2e^{\dagger}_{-2,0}-3e^{\dagger}_{-1,0}+e^{\dagger}_{1,0})\mathcal{Q}_{-1}\otimes\mathcal{H}_{-1}+(e^{\dagger}_{-2,0}-3e^{\dagger}_{-1,0}-2e^{\dagger}_{1,-1}-e^{\dagger}_{1,0})\mathcal{Q}_0\otimes\mathcal{H}_{-2}\\
&+\beta_{-2,0}\mathcal{H}_{-2}\otimes\mathcal{H}_0-(2\beta_{-2,0}+\beta_{1,0}-3\beta_{-1,0})\mathcal{H}_{-1}\otimes\mathcal{H}_{-1}+(\beta_{-2,0}-2\beta_{1,-1}-3\beta_{-1,0}-\beta_{1,0})\mathcal{H}_0\otimes\mathcal{H}_{-2}\\
&+\mu_{-2,0}\mathcal{Q}_{-2}\otimes\mathcal{Q}_0-(2\mu_{-2,0}+\mu_{1,0}-3\mu_{-1,0})\mathcal{Q}_{-1}\otimes\mathcal{Q}_{-1}+(\mu_{-2,0}-2\mu_{1,-1}-3\mu_{-1,0}-\mu_{1,0})\mathcal{Q}_0\otimes\mathcal{Q}_{-2}.
\end{aligned}$$

Applying d_0 to $[\mathcal{L}_2,\mathcal{L}_{-2}]=4\mathcal{L}_0$, we have

$$\mathcal{L}_2\cdot d_0(\mathcal{L}_{-2})=\mathcal{L}_{-2}\cdot d_0(\mathcal{L}_2). \tag{3.6}$$

Comparing the coefficients of $\mathcal{L}_2\otimes\mathcal{H}_{-2}$, $\mathcal{L}_{-2}\otimes\mathcal{H}_2$, $\mathcal{L}_2\otimes\mathcal{Q}_{-2}$, $\mathcal{L}_{-2}\otimes\mathcal{Q}_2$, $\mathcal{G}_2\otimes\mathcal{H}_{-2}$, $\mathcal{G}_{-2}\otimes\mathcal{H}_2$, $\mathcal{G}_2\otimes\mathcal{Q}_{-2}$, $\mathcal{G}_{-2}\otimes\mathcal{Q}_2$ in (3.6), one has $a_{2,0}=a_{-2,0}=c_{2,0}=c_{-2,0}=d^{\dagger}_{2,0}=d^{\dagger}_{-2,0}=f_{2,0}=f_{-2,0}=0$. Comparing the coefficients of $\mathcal{H}_{-2}\otimes\mathcal{L}_2$, $\mathcal{H}_2\otimes\mathcal{L}_{-2}$, $\mathcal{Q}_{-2}\otimes\mathcal{L}_2$, $\mathcal{Q}_2\otimes\mathcal{L}_{-2}$, $\mathcal{H}_{-2}\otimes\mathcal{G}_2$, $\mathcal{H}_2\otimes\mathcal{G}_{-2}$, $\mathcal{Q}_{-2}\otimes\mathcal{G}_2$, $\mathcal{Q}_2\otimes\mathcal{G}_{-2}$ in (3.6), one has $a^{\dagger}_{-2,0}=a^{\dagger}_{2,0}=c^{\dagger}_{-2,0}=c^{\dagger}_{2,0}=d_{-2,0}=d_{2,0}=f^{\dagger}_{-2,0}=f^{\dagger}_{2,0}=0$. Comparing the coefficients of $\mathcal{L}_0\otimes\mathcal{G}_0$, $\mathcal{G}_0\otimes\mathcal{L}_0$, $\mathcal{L}_0\otimes\mathcal{L}_0$ and $\mathcal{G}_0\otimes\mathcal{G}_0$ in (3.6), one has $b_{-2,0}=b^{\dagger}_{-2,0}=\alpha_{2,0}=k_{2,0}=0$. Comparing the coefficients of $\mathcal{H}_1\otimes\mathcal{Q}_{-1}$, $\mathcal{Q}_1\otimes\mathcal{H}_{-1}$, $\mathcal{H}_0\otimes\mathcal{H}_0$ and $\mathcal{Q}_0\otimes\mathcal{Q}_0$ in (3.6), one has

$$x_{-2,0}+x_{2,0}=x_{-1,0}+x_{1,0},\ \ y_{-2,0}+y_{2,0}=y_{-1,0}+y_{1,0}\ \ \text{for } x=e,\beta,\ \ y=e^{\dagger},\mu. \tag{3.7}$$

Thus one can rewrite $d_0(\mathcal{L}_1)-\mu_{1,-1}\mathcal{Q}_0\otimes\mathcal{Q}_1-\mu_{1,0}\mathcal{Q}_1\otimes\mathcal{Q}_0$ and $d_0(\mathcal{L}_{-1})$ respectively as

$$\begin{aligned}
&e_{1,-1}\mathcal{H}_0\otimes\mathcal{Q}_1+e_{1,0}\mathcal{H}_1\otimes\mathcal{Q}_0+e^{\dagger}_{1,-1}\mathcal{Q}_0\otimes\mathcal{H}_1+e^{\dagger}_{1,0}\mathcal{Q}_1\otimes\mathcal{H}_0+j_{1,-1}\mathcal{H}_0\otimes\mathcal{H}_1+j_{1,0}\mathcal{H}_1\otimes\mathcal{H}_0,\\
&e_{-1,0}\mathcal{H}_{-1}\otimes\mathcal{Q}_0-(e_{-1,0}+e_{1,-1}+e_{1,0})\mathcal{H}_0\otimes\mathcal{Q}_{-1}+e^{\dagger}_{-1,0}\mathcal{Q}_{-1}\otimes\mathcal{H}_0-(e^{\dagger}_{-1,0}+e^{\dagger}_{1,-1}+e^{\dagger}_{1,0})\mathcal{Q}_0\otimes\mathcal{H}_{-1}\\
&+\beta_{-1,0}\mathcal{H}_{-1}\otimes\mathcal{H}_0-(\beta_{-1,0}+\beta_{1,-1}+\beta_{1,0})\mathcal{H}_0\otimes\mathcal{H}_{-1}+\mu_{-1,0}\mathcal{Q}_{-1}\otimes\mathcal{Q}_0-(\mu_{-1,0}+\mu_{1,-1}+\mu_{1,0})\mathcal{Q}_0\otimes\mathcal{Q}_{-1}.
\end{aligned}$$

One can write

$$\begin{aligned}
d_0(\mathcal{G}_n)=\sum_{i\in\mathbb{Z}}(&m_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{H}_{-i}+m^{\dagger}_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{L}_{-i}+k_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{G}_{-i}+k^{\dagger}_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{L}_{-i}\\
&+u_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{Q}_{-i}+u^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{L}_{-i}+p_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{G}_{-i}+p^{\dagger}_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{H}_{-i}\\
&+q_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{Q}_{-i}+q^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{H}_{-i}+s_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{Q}_{-i}+s^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{G}_{-i}\\
&+t_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{L}_{-i}+w_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{H}_{-i}+t^{\dagger}_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{G}_{-i}+w^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{Q}_{-i}).
\end{aligned}$$

Applying d_0 to $[\mathcal{L}_1, \mathcal{G}_1] = 0$, we have $\mathcal{L}_1 \cdot d_0(\mathcal{G}_1) = (-1)^{[d]}\mathcal{G}_1 \cdot d_0(\mathcal{L}_1)$. Comparing the coefficients of $\mathcal{L}_{i+1} \otimes \mathcal{H}_{1-i}$, one has $(i-1)m_{1,i-1} = im_{1,i}$ for $i \neq 0, 1$, which together with $\{i \,|\, m_{1,i} \neq 0\}$ being finite, implies $m_{1,i} = e^{\dagger}_{1,-1} = e^{\dagger}_{1,0} = 0,\ \forall\ i \in \mathbb{Z}^*$. Using the same techniques to $\mathcal{L}_{i+1} \otimes \mathcal{Q}_{1-i}$, $\mathcal{G}_{i+1} \otimes \mathcal{H}_{1-i}$, $\mathcal{G}_{i+1} \otimes \mathcal{Q}_{1-i}$, one has

$$x_{1,i} = y_{1,-1} = y_{1,0} = 0, \text{ for } x = u, p^{\dagger}, s,\ y = \mu, \beta, e,\ \forall\ i \in \mathbb{Z}^*.$$

Comparing the coefficients of $\mathcal{H}_{i+1} \otimes \mathcal{L}_{1-i}$, $\mathcal{Q}_{i+1} \otimes \mathcal{L}_{1-i}$, $\mathcal{H}_{i+1} \otimes \mathcal{G}_{1-i}$, $\mathcal{Q}_{i+1} \otimes \mathcal{G}_{1-i}$, one has $m^{\dagger}_{1,i} = u^{\dagger}_{1,i} = p_{1,i} = s^{\dagger}_{1,i} = 0,\ \forall\ i \neq -1$. Comparing the coefficients of $\mathcal{L}_{i+1}\otimes\mathcal{G}_{1-i}$, $\mathcal{G}_{i+1}\otimes\mathcal{L}_{1-i}$, $\mathcal{L}_{i+1} \otimes \mathcal{L}_{1-i}$, $\mathcal{G}_{i+1} \otimes \mathcal{G}_{1-i}$, one has $x_{1,i} = 0,\ x_{1,-1} = -x_{1,0}$ for $x = k, t, k^{\dagger}, t^{\dagger},\ \forall\ i \neq 0, -1$. Comparing the coefficients of $\mathcal{H}_{i+1} \otimes \mathcal{Q}_{1-i}$, $\mathcal{Q}_{i+1} \otimes \mathcal{H}_{1-i}$, $\mathcal{H}_{i+1} \otimes \mathcal{H}_{1-i}$, $\mathcal{Q}_{i+1} \otimes \mathcal{Q}_{1-i}$, one has $q_{1,i} = q^{\dagger}_{1,i} = w_{1,i} = w^{\dagger}_{1,i} = 0$ for $i \in \mathbb{Z}$. Also by the above results, we have $d_0(\mathcal{L}_1) = 0$.

Applying d_0 to $[\mathcal{L}_1, \mathcal{G}_0] = \mathcal{G}_1$, we have $\mathcal{L}_1 \cdot d_0(\mathcal{G}_0) = d_0(\mathcal{G}_1)$. Comparing the coefficients of $\mathcal{L}_{i+1} \otimes \mathcal{H}_{-i}$, one has $(i-1)m_{0,i} = (i+1)m_{0,i+1}$ for $i \neq 0$. Since $\{i \mid\ m_{0,i} \neq 0\}$ is a finite set, we obtain $m_{0,i} = 0,\ m_{0,1} = -m_{0,0} + m_{1,0}$ for $i \neq 0, 1$. Applying the same techniques to $\mathcal{L}_{i+1} \otimes \mathcal{Q}_{-i}$, $\mathcal{G}_{i+1} \otimes \mathcal{H}_{-i}$, $\mathcal{G}_{i+1} \otimes \mathcal{Q}_{-i}$, $\mathcal{H}_i \otimes \mathcal{L}_{1-i}$, $\mathcal{Q}_i \otimes \mathcal{L}_{1-i}$, $\mathcal{H}_i \otimes \mathcal{G}_{1-i}$, $\mathcal{Q}_i \otimes \mathcal{G}_{1-i}$, we have $x_{0,i} = 0,\ x_{0,1} = -x_{0,0}+x_{1,0}$ and $y_{0,j} = 0,\ y_{0,-1} = -y_{0,0}+y_{1,-1}$ for $x = u, p^{\dagger}, s, y = m^{\dagger}, u^{\dagger}, p, s^{\dagger}$, $\forall\ i \neq 0, 1, j \neq 0, -1$. Comparing the coefficients of $\mathcal{L}_{i+1} \otimes \mathcal{G}_{-i}$, $\mathcal{G}_{i+1} \otimes \mathcal{L}_{-i}$, $\mathcal{L}_{i+1} \otimes \mathcal{L}_{-i}$, $\mathcal{G}_{i+1} \otimes \mathcal{G}_{-i}$, we have

$$x_{0,i} = x_{0,1} + \frac{1}{2}x_{0,0} - \frac{1}{2}x_{1,0} = x_{0,-1} + \frac{1}{2}x_{0,0} + \frac{1}{2}x_{1,0} = 0 \text{ for } x = k, k^{\dagger}, t, t^{\dagger},\ \forall\ i \neq 0, -1, 1.$$

Comparing the coefficients of $\mathcal{H}_i \otimes \mathcal{Q}_{1-i}$, $\mathcal{Q}_i \otimes \mathcal{H}_{1-i}$, $\mathcal{H}_i \otimes \mathcal{H}_{1-i}$, $\mathcal{Q}_i \otimes \mathcal{Q}_{1-i}$, one has

$$q_{0,i} = q^{\dagger}_{0,i} = w_{0,i} = w^{\dagger}_{0,i} = 0,\ \forall\ i \neq 0.$$

Applying d_0 to $[\mathcal{L}_{-1}, \mathcal{G}_1] = -2\mathcal{G}_0$, we have $\mathcal{L}_{-1} \cdot d_0(\mathcal{G}_1) + 2d_0(\mathcal{G}_0) = (-1)^{[d]}\mathcal{G}_1 \cdot d_0(\mathcal{L}_{-1})$. Comparing the coefficients of $\mathcal{L}_0 \otimes \mathcal{H}_0$, $\mathcal{H}_0 \otimes \mathcal{L}_0$, $\mathcal{L}_0 \otimes \mathcal{Q}_0$, $\mathcal{Q}_0 \otimes \mathcal{L}_0$, one has

$$\begin{aligned} m_{1,0} &= m_{0,0} - (-1)^{[d]}e^{\dagger}_{-1,0},\ m^{\dagger}_{1,-1} = m^{\dagger}_{0,0} + (-1)^{[d]}e_{-1,0}, \\ u_{1,0} &= -u_{0,0} + (-1)^{[d]}\mu_{-1,0},\ u^{\dagger}_{1,-1} = u^{\dagger}_{0,0} + (-1)^{[d]}\mu_{-1,0}. \end{aligned}$$

Comparing the coefficients of $\mathcal{H}_0 \otimes \mathcal{G}_0$, $\mathcal{G}_0 \otimes \mathcal{H}_0$, $\mathcal{G}_0 \otimes \mathcal{Q}_0$, $\mathcal{Q}_0 \otimes \mathcal{G}_0$, $\mathcal{L}_0 \otimes \mathcal{G}_0$, $\mathcal{G}_0 \otimes \mathcal{L}_0$, $\mathcal{L}_0 \otimes \mathcal{L}_0$, $\mathcal{G}_0 \otimes \mathcal{G}_0$, we have $y_{0,0} = y^{\dagger}_{0,0} = t_{0,0} = t^{\dagger}_{0,0} = 0$, $p^{\dagger}_{1,0} = p^{\dagger}_{0,0} + (-1)^{[d]}\frac{1}{2}\beta_{-1,0}$, $p_{1,-1} = p_{0,0} + (-1)^{[d]+1}\frac{1}{2}\beta_{-1,0}$ and $s_{1,0} = s_{0,0} + (-1)^{[d]}\frac{1}{2}e_{-1,0}$, $s^{\dagger}_{1,-1} = s^{\dagger}_{0,0} + (-1)^{[d]}\frac{1}{2}e^{\dagger}_{-1,0}$. Comparing the coefficients of $\mathcal{H}_0 \otimes \mathcal{Q}_0$, $\mathcal{Q}_0 \otimes \mathcal{H}_0$, $\mathcal{H}_0 \otimes \mathcal{H}_0$, $\mathcal{Q}_0 \otimes \mathcal{Q}_0$, one has

$$q^{\dagger}_{0,0} = q_{0,0} = (-1)^{[d]}\mu_{-1,0},\quad w_{0,0} = (-1)^{[d]+1}2e_{-1,0} + (-1)^{[d]}2e^{\dagger}_{-1,0},\quad w^{\dagger}_{0,0} = 0.$$

Applying d_0 to $[\mathcal{L}_{-1}, \mathcal{G}_0] = -\mathcal{G}_{-1}$, one has $\mathcal{L}_{-1} \cdot d_0(\mathcal{G}_0) + d_0(\mathcal{G}_{-1}) = (-1)^{[d]}\mathcal{G}_0 \cdot d_0(\mathcal{L}_{-1})$. Comparing the coefficients of $\mathcal{L}_{-1} \otimes \mathcal{H}_0$, $\mathcal{L}_0 \otimes \mathcal{H}_{-1}$, $\mathcal{L}_1 \otimes \mathcal{H}_{-2}$, $\mathcal{H}_0 \otimes \mathcal{L}_{-1}$, $\mathcal{H}_{-2} \otimes \mathcal{L}_1$, $\mathcal{H}_{-1} \otimes \mathcal{L}_0$, we have $m_{-1,0} = m_{0,0} + (-1)^{[d]}2e^{\dagger}_{-1,0},\ m_{-1,1} = (-1)^{[d]+1}4e^{\dagger}_{-1,0},\ m_{-1,2} = (-1)^{[d]}e^{\dagger}_{-1,0},$

$$m^{\dagger}_{-1,1} = m^{\dagger}_{0,0} + (-1)^{[d]+1}2e_{-1,0},\ m^{\dagger}_{-1,-1} = (-1)^{[d]+1}e_{-1,0},\ m^{\dagger}_{-1,0} = (-1)^{[d]}4e_{-1,0}.$$

According to $\mathcal{L}_{-1} \otimes \mathcal{Q}_0$, $\mathcal{L}_0 \otimes \mathcal{Q}_{-1}$, $\mathcal{L}_1 \otimes \mathcal{Q}_{-2}$, $\mathcal{Q}_0 \otimes \mathcal{L}_{-1}$, $\mathcal{Q}_{-2} \otimes \mathcal{L}_1$, $\mathcal{Q}_{-1} \otimes \mathcal{L}_0$, one has

$$u_{-1,0} = u_{0,0} + (-1)^{[d]}2\mu_{-1,0},\ u_{-1,1} = (-1)^{[d]+1}4\mu_{-1,0},\ u_{-1,2} = (-1)^{[d]}\mu_{-1,0},$$
$$u^{\dagger}_{-1,1} = u^{\dagger}_{0,0} + (-1)^{[d]}2\mu_{-1,0},\ u^{\dagger}_{-1,-1} = (-1)^{[d]}\mu_{-1,0},\ u^{\dagger}_{-1,0} = (-1)^{[d]+1}4\mu_{-1,0}.$$

Comparing the coefficients of $\mathcal{G}_{-1} \otimes \mathcal{H}_0$, $\mathcal{G}_0 \otimes \mathcal{H}_{-1}$, $\mathcal{G}_1 \otimes \mathcal{H}_{-2}$, $\mathcal{H}_0 \otimes \mathcal{G}_{-1}$, $\mathcal{H}_{-2} \otimes \mathcal{G}_1$, $\mathcal{H}_{-1} \otimes \mathcal{G}_0$, we obtain $p^{\dagger}_{-1,0} = p^{\dagger}_{0,0} + (-1)^{[d]+1}\beta_{-1,0},\ p^{\dagger}_{-1,1} = (-1)^{[d]}2\beta_{-1,0},\ p^{\dagger}_{-1,2} = (-1)^{[d]+1}0.5\beta_{-1,0},$

$$p_{-1,1} = p_{0,0} + (-1)^{[d]}\beta_{-1,0},\ p_{-1,-1} = (-1)^{[d]}0.5\beta_{-1,0},\ p_{-1,0} = (-1)^{[d]+1}2\beta_{-1,0}.$$

Comparing the coefficients of $\mathcal{G}_{-1} \otimes \mathcal{Q}_0$, $\mathcal{G}_0 \otimes \mathcal{Q}_{-1}$, $\mathcal{G}_1 \otimes \mathcal{Q}_{-2}$, $\mathcal{Q}_0 \otimes \mathcal{G}_{-1}$, $\mathcal{Q}_{-2} \otimes \mathcal{G}_1$, $\mathcal{Q}_{-1} \otimes \mathcal{G}_0$, one has $s_{-1,0} = s_{0,0} + (-1)^{[d]+1}e_{-1,0},\ s_{-1,1} = (-1)^{[d]}2e_{-1,0},\ s_{-1,2} = (-1)^{[d]+1}0.5e_{-1,0},$

$$s^{\dagger}_{-1,1} = s^{\dagger}_{0,0} + (-1)^{[d]+1}e^{\dagger}_{-1,0},\ s^{\dagger}_{-1,-1} = (-1)^{[d]+1}0.5e^{\dagger}_{-1,0},\ s^{\dagger}_{-1,2} = (-1)^{[d]}2e^{\dagger}_{-1,0}.$$

Comparing the coefficients of $\mathcal{L}_0 \otimes \mathcal{G}_{-1}$, $\mathcal{L}_{-2} \otimes \mathcal{G}_1$, $\mathcal{L}_{-1} \otimes \mathcal{G}_0$, $\mathcal{G}_{-1} \otimes \mathcal{L}_0$, $\mathcal{G}_0 \otimes \mathcal{L}_{-1}$, $\mathcal{G}_{-2} \otimes \mathcal{L}_1$, $\mathcal{L}_{-1} \otimes \mathcal{L}_0$, $\mathcal{L}_0 \otimes \mathcal{L}_{-1}$, $\mathcal{L}_{-2} \otimes \mathcal{L}_1$, $\mathcal{L}_1 \otimes \mathcal{L}_{-2}$, $\mathcal{G}_{-1} \otimes \mathcal{G}_0$, $\mathcal{G}_0 \otimes \mathcal{G}_{-1}$, $\mathcal{G}_{-2} \otimes \mathcal{G}_1$, $\mathcal{G}_1 \otimes \mathcal{G}_{-2}$, we have

$$x_{-1,0} + x_{1,0} = x_{-1,1} - x_{1,0} = x_{-1,-1} = x_{-1,2} = 0 \ \text{ for } x = k, k^{\dagger}, t, t^{\dagger}.$$

Comparing the coefficients of $\mathcal{H}_{-1} \otimes \mathcal{Q}_0$, $\mathcal{Q}_0 \otimes \mathcal{H}_{-1}$, $\mathcal{H}_0 \otimes \mathcal{H}_{-1}$, $\mathcal{H}_{-1} \otimes \mathcal{H}_0$, one has

$$q_{-1,0} = (-1)^{[d]}2\mu_{-1,0},\ q^{\dagger}_{-1,1} = (-1)^{[d]}2\mu_{-1,0},\ w_{-1,1} = (-1)^{[d]+1}2e_{-1,0},\ w_{-1,0} = (-1)^{[d]}2e^{\dagger}_{-1,0}.$$

Applying d_0 to $[\mathcal{L}_2, \mathcal{G}_{-1}] = 3\mathcal{G}_1$, we have $\mathcal{L}_2 \cdot d_0(\mathcal{G}_{-1}) = (-1)^{[d]}\mathcal{G}_{-1} \cdot d_0(\mathcal{L}_2) + 3d_0(\mathcal{G}_1)$. Comparing the coefficients of $\mathcal{L}_2 \otimes \mathcal{H}_{-1}$, $\mathcal{L}_0 \otimes \mathcal{H}_1$, $\mathcal{H}_{-1} \otimes \mathcal{L}_2$, $\mathcal{H}_1 \otimes \mathcal{L}_0$, $\mathcal{L}_2 \otimes \mathcal{Q}_{-1}$, $\mathcal{L}_0 \otimes \mathcal{Q}_1$, $\mathcal{G}_2 \otimes \mathcal{H}_{-1}$, $\mathcal{G}_0 \otimes \mathcal{H}_1$, $\mathcal{L}_1 \otimes \mathcal{G}_0$, $\mathcal{G}_2 \otimes \mathcal{L}_{-1}$, $\mathcal{L}_0 \otimes \mathcal{L}_1$, $\mathcal{G}_0 \otimes \mathcal{G}_1$ and combine with (3.7), we have $\theta_{-1,0} = \theta_{-2,0} = \theta_{2,0} = x_{1,0} = y_{1,0} - y_{0,0} = z_{1,-1} - z_{0,0} = 0$ for $\theta = e^{\dagger}, e, \mu, \beta, x = k, k^{\dagger}, t, t^{\dagger}, y = m, u, p^{\dagger}, s,\ z = m^{\dagger}, u^{\dagger}, s^{\dagger}, p$. Thus $d_0(\mathcal{L}_{-1}) = d_0(\mathcal{L}_{-2}) = d_0(\mathcal{L}_2) = 0$.

One can write

$$\begin{aligned} d_0(\mathcal{H}_n) = \sum_{i\in\mathbb{Z}} (&\xi_{n,i}\mathcal{L}_{i+n} \otimes \mathcal{H}_{-i} + \xi^{\dagger}_{n,i}\mathcal{H}_{i+n} \otimes \mathcal{L}_{-i} + \eta_{n,i}\mathcal{L}_{i+n} \otimes \mathcal{G}_{-i} + \eta^{\dagger}_{n,i}\mathcal{G}_{i+n} \otimes \mathcal{L}_{-i} \\ &+\varepsilon_{n,i}\mathcal{L}_{i+n} \otimes \mathcal{Q}_{-i} + \varepsilon^{\dagger}_{n,i}\mathcal{Q}_{i+n} \otimes \mathcal{L}_{-i} + \delta_{n,i}\mathcal{H}_{i+n} \otimes \mathcal{G}_{-i} + \delta^{\dagger}_{n,i}\mathcal{G}_{i+n} \otimes \mathcal{H}_{-i} \\ &+\psi_{n,i}\mathcal{H}_{i+n} \otimes \mathcal{Q}_{-i} + \psi^{\dagger}_{n,i}\mathcal{Q}_{i+n} \otimes \mathcal{H}_{-i} + \varphi_{n,i}\mathcal{G}_{i+n} \otimes \mathcal{Q}_{-i} + \varphi^{\dagger}_{n,i}\mathcal{Q}_{i+n} \otimes \mathcal{G}_{-i} \\ &+h_{n,i}\mathcal{L}_{i+n} \otimes \mathcal{L}_{-i} + l_{n,i}\mathcal{H}_{i+n} \otimes \mathcal{H}_{-i} + g_{n,i}\mathcal{G}_{i+n} \otimes \mathcal{G}_{-i} + v_{n,i}\mathcal{Q}_{i+n} \otimes \mathcal{Q}_{-i}). \end{aligned}$$

Applying d_0 to $[\mathcal{L}_1, \mathcal{H}_0] = 0$, we have $\mathcal{L}_1 \cdot d_0(\mathcal{H}_0) = 0$. Comparing the coefficients of $\mathcal{L}_{i+1} \otimes \mathcal{H}_{-i}$, we have $(i-1)\xi_{0,i} = (i+1)\xi_{0,i+1}$ for $i \in \mathbb{Z}$, which implies $\xi_{0,i} = 0$, $\xi_{0,1} = -\xi_{0,0}$,

$\forall\ i\neq 0,1$. Comparing the coefficients of $\mathcal{L}_{i+1}\otimes\mathcal{Q}_{-i}$, $\mathcal{G}_{i+1}\otimes\mathcal{H}_{-i}$, $\mathcal{G}_{i+1}\otimes\mathcal{Q}_{-i}$, $\mathcal{H}_{i+1}\otimes\mathcal{L}_{-i}$, $\mathcal{Q}_{i+1}\otimes\mathcal{L}_{-i}$, $\mathcal{H}_{i+1}\otimes\mathcal{G}_{-i}$, $\mathcal{Q}_{i+1}\otimes\mathcal{G}_{-i}$, we have

$$x_{0,i}=y_{0,j}=x_{0,1}+x_{0,0}=y_{0,-1}+y_{0,0}=0,\ \ x=\varepsilon,\delta^{\dagger},\varphi,\ \ y=\xi^{\dagger},\varepsilon^{\dagger},\delta,\varphi^{\dagger},\forall\ i\neq 0,1,j\neq 0,-1.$$

Comparing the coefficients of $\mathcal{L}_{i+1}\otimes\mathcal{G}_{-i}$, $\mathcal{G}_{i+1}\otimes\mathcal{L}_{-i}$, $\mathcal{L}_{i+1}\otimes\mathcal{L}_{-i}$, $\mathcal{G}_{i+1}\otimes\mathcal{G}_{-i}$, we have

$$x_{0,i}=x_{0,-1}+\frac{1}{2}x_{0,0}=x_{0,1}-\frac{1}{2}x_{0,0}=0\ \text{for}\,x=\eta,\eta^{\dagger},h,g,\forall\ i\neq 0,1,-1.$$

Comparing the coefficients of $\mathcal{H}_{i+1}\otimes\mathcal{Q}_{-i}$, $\mathcal{Q}_{i+1}\otimes\mathcal{H}_{-i}$, $\mathcal{H}_{i+1}\otimes\mathcal{H}_{-i}$, $\mathcal{Q}_{i+1}\otimes\mathcal{Q}_{-i}$, one has

$$x_{0,i}=0\ \ \text{for}\ \ x=\psi,\psi^{\dagger},l,v,\forall\ i\neq 0.$$

Applying d_0 to $[\mathcal{H}_0,\mathcal{G}_0]=\mathcal{G}_0$, one has $\mathcal{H}_0\cdot d_0(\mathcal{G}_0)=(-1)^{[d]}\mathcal{G}_0\cdot d_0(\mathcal{H}_0)+d_0(\mathcal{G}_0)$. Comparing the coefficients of $\mathcal{L}_0\otimes\mathcal{Q}_0$, $\mathcal{Q}_0\otimes\mathcal{L}_0$, $\mathcal{G}_0\otimes\mathcal{H}_0$, $\mathcal{G}_1\otimes\mathcal{H}_{-1}$, $\mathcal{H}_{-1}\otimes\mathcal{G}_1$, $\mathcal{G}_0\otimes\mathcal{L}_0$, $\mathcal{G}_0\otimes\mathcal{G}_0$, $\mathcal{G}_1\otimes\mathcal{G}_{-1}$, $\mathcal{L}_1\otimes\mathcal{L}_{-1}$, $\mathcal{L}_{-1}\otimes\mathcal{L}_1$, $\mathcal{G}_0\otimes\mathcal{Q}_0$, $\mathcal{Q}_0\otimes\mathcal{G}_0$, $\mathcal{L}_0\otimes\mathcal{H}_0$, $\mathcal{H}_0\otimes\mathcal{L}_0$, one has

$$\begin{aligned}&l_{0,0}=\varepsilon_{0,0}=\varepsilon^{\dagger}_{0,0}=0,u^{\dagger}_{0,0}=-u_{0,0},\delta^{\dagger}_{0,0}=\delta_{0,0},\xi_{0,0}=\xi^{\dagger}_{0,0}=-2\varphi_{0,0},\varphi^{\dagger}_{0,0}=-\varphi_{0,0},h_{0,0}=-4\varphi_{0,0},\\&v_{0,0}=(-1)^{[d]+1}u_{0,0},\psi_{0,0}=(-1)^{[d]}s_{0,0},\psi^{\dagger}_{0,0}=(-1)^{[d]+1}s^{\dagger}_{0,0},m_{0,0}=2s^{\dagger}_{0,0},\ m^{\dagger}_{0,0}=-2s_{0,0}.\end{aligned}$$

Denoting $v_1=\mathcal{Q}_0\otimes\mathcal{H}_0$, $v_2=\mathcal{H}_0\otimes\mathcal{Q}_0$ and observing the facts

$$\mathcal{G}_0\cdot v_1=2\mathcal{L}_0\otimes\mathcal{H}_0+\mathcal{Q}_0\otimes\mathcal{G}_0,\ \mathcal{G}_0\cdot v_2=-\mathcal{G}_0\otimes\mathcal{Q}_0+2\mathcal{H}_0\otimes\mathcal{L}_0,\ \mathcal{L}_{\pm1}\cdot v_i=\mathcal{L}_{\pm2}\cdot v_i=0,$$

for $i=1,2$, replacing d_0 by $d_0-\kappa_{\rm inn}$ (where κ is some linear combination of v_1, v_2), one has

$$\begin{aligned}d_0(\mathcal{G}_0)&=p^{\dagger}_{0,0}\mathcal{G}_0\otimes\mathcal{H}_0+p_{0,0}\mathcal{H}_0\otimes\mathcal{G}_0+u_{0,0}\mathcal{L}_0\otimes\mathcal{Q}_0-u_{0,0}\mathcal{Q}_0\otimes\mathcal{L}_0,\\d_0(\mathcal{G}_1)&=-u_{0,0}\mathcal{Q}_0\otimes\mathcal{L}_1+u_{0,0}\mathcal{L}_1\otimes\mathcal{Q}_0+p_{0,0}\mathcal{H}_0\otimes\mathcal{G}_1+p^{\dagger}_{0,0}\mathcal{G}_1\otimes\mathcal{H}_0,\\d_0(\mathcal{G}_{-1})&=u_{0,0}\mathcal{L}_{-1}\otimes\mathcal{Q}_0-u_{0,0}\mathcal{Q}_0\otimes\mathcal{L}_{-1}+p^{\dagger}_{0,0}\mathcal{G}_{-1}\otimes\mathcal{H}_0+p_{0,0}\mathcal{H}_0\otimes\mathcal{G}_{-1}.\end{aligned}$$

Comparing the coefficients of $\mathcal{G}_0\otimes\mathcal{H}_{-1}$, $\mathcal{G}_{-2}\otimes\mathcal{G}_1$ in $\mathcal{H}_0\cdot d_0(\mathcal{G}_{-1})=(-1)^{[d]}\mathcal{G}_{-1}\cdot d_0(\mathcal{H}_0)+d_0(\mathcal{G}_{-1})$, we have $\varphi_{0,0}=\delta_{0,0}=0$. Comparing the coefficients of $\mathcal{L}_{-1}\otimes\mathcal{G}_0$, $\mathcal{G}_{-1}\otimes\mathcal{L}_0$, $\mathcal{G}_{-1}\otimes\mathcal{G}_0$ in $\mathcal{L}_{-1}\cdot d_0(\mathcal{H}_0)=\mathcal{H}_0\cdot d_0(\mathcal{L}_{-1})$, we have $\eta_{0,0}=\eta^{\dagger}_{0,0}=g_{0,0}=0$. Thus one can suppose

$$d_0(\mathcal{H}_0)=(-1)^{[d]+1}u_{0,0}\mathcal{Q}_0\otimes\mathcal{Q}_0.$$

Write

$$\begin{aligned}d_0(\mathcal{Q}_n)=\sum_{i\in\mathbb{Z}}\big(&A_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{H}_{-i}+A^{\dagger}_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{L}_{-i}+B_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{G}_{-i}+B^{\dagger}_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{L}_{-i}\\&+D_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{Q}_{-i}+D^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{L}_{-i}+E_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{G}_{-i}+E^{\dagger}_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{H}_{-i}\\&+F_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{Q}_{-i}+F^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{H}_{-i}+R_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{Q}_{-i}+R^{\dagger}_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{G}_{-i}\\&+X_{n,i}\mathcal{L}_{i+n}\otimes\mathcal{L}_{-i}+Y_{n,i}\mathcal{H}_{i+n}\otimes\mathcal{H}_{-i}+M_{n,i}\mathcal{G}_{i+n}\otimes\mathcal{G}_{-i}+N_{n,i}\mathcal{Q}_{i+n}\otimes\mathcal{Q}_{-i}\big).\end{aligned}$$

Comparing the coefficients of $\mathcal{G}_0\otimes\mathcal{H}_0$, $\mathcal{H}_0\otimes\mathcal{G}_0$, $\mathcal{L}_0\otimes\mathcal{G}_0$, $\mathcal{G}_0\otimes\mathcal{L}_0$, $\mathcal{G}_0\otimes\mathcal{G}_0$, $\mathcal{Q}_0\otimes\mathcal{Q}_0$, $\mathcal{L}_0\otimes\mathcal{H}_0$, $\mathcal{H}_0\otimes\mathcal{L}_0$, $\mathcal{Q}_0\otimes\mathcal{G}_0$, $\mathcal{G}_0\otimes\mathcal{Q}_0$, $\mathcal{L}_0\otimes\mathcal{L}_0$, $\mathcal{H}_0\otimes\mathcal{H}_0$ in $\mathcal{H}_0\cdot d_0(\mathcal{Q}_0)+d_0(\mathcal{Q}_0)=(-1)^{[d]}\mathcal{Q}_0\cdot d_0(\mathcal{H}_0)$, we obtain $E^{\dagger}_{0,0}=E_{0,0}=B_{0,0}=B^{\dagger}_{0,0}=M_{0,0}=X_{0,0}=Y_{0,0}=A_{0,0}=A^{\dagger}_{0,0}=R^{\dagger}_{0,0}=R_{0,0}=N_{0,0}=0$. Thus we can rewrite $d_0(\mathcal{Q}_0)$ as

$$D_{0,0}\mathcal{L}_0\otimes\mathcal{Q}_0-D_{0,0}\mathcal{L}_1\otimes\mathcal{Q}_{-1}+D^{\dagger}_{0,0}\mathcal{Q}_0\otimes\mathcal{L}_0-D^{\dagger}_{0,0}\mathcal{Q}_{-1}\otimes\mathcal{L}_1+F_{0,0}\mathcal{H}_0\otimes\mathcal{Q}_0+F^{\dagger}_{0,0}\mathcal{Q}_0\otimes\mathcal{H}_0.$$

Applying d_0 to $[\mathcal{G}_0,\mathcal{Q}_0]=2\mathcal{L}_0$, we have $\mathcal{G}_0\cdot d_0(\mathcal{Q}_0)+\mathcal{Q}_0\cdot d_0(\mathcal{G}_0)=0$. Comparing the coefficients of $\mathcal{G}_0\otimes\mathcal{Q}_0$, $\mathcal{Q}_0\otimes\mathcal{G}_0$, $\mathcal{H}_0\otimes\mathcal{L}_0$, $\mathcal{L}_0\otimes\mathcal{H}_0$, $\mathcal{G}_1\otimes\mathcal{Q}_{-1}$, $\mathcal{H}_{-1}\otimes\mathcal{L}_1$, we have $F_{0,0}=F^{\dagger}_{0,0}=-p_{0,0},\ p^{\dagger}_{0,0}=p_{0,0}, D_{0,0}=D^{\dagger}_{0,0}=0$. Thus one can suppose

$$\begin{aligned}d_0(\mathcal{Q}_0)&=-p_{0,0}\mathcal{H}_0\otimes\mathcal{Q}_0-p_{0,0}\mathcal{Q}_0\otimes\mathcal{H}_0,\\ d_0(\mathcal{G}_0)&=p_{0,0}\mathcal{G}_0\otimes\mathcal{H}_0+p_{0,0}\mathcal{H}_0\otimes\mathcal{G}_0+u_{0,0}\mathcal{L}_0\otimes\mathcal{Q}_0-u_{0,0}\mathcal{Q}_0\otimes\mathcal{L}_0.\end{aligned}$$

Denoting $x=\mathcal{H}_0\otimes\mathcal{H}_0$, observing $\mathcal{G}_0\cdot x=-\mathcal{G}_0\otimes\mathcal{H}_0-\mathcal{H}_0\otimes\mathcal{G}_0$ and replacing d_0 by $d_0+p_{0,0}x_{\rm inn}$ (noting that this replacement does not affect $d_0(\mathcal{L}_{\pm1}),\ d_0(\mathcal{L}_{\pm2})$), one can suppose

$$\begin{aligned}&d_0(\mathcal{Q}_0)=0,\ d_0(\mathcal{H}_0)=(-1)^{[d]+1}u_{0,0}\mathcal{Q}_0\otimes\mathcal{Q}_0,\ d_0(\mathcal{G}_0)=u_{0,0}\mathcal{L}_0\otimes\mathcal{Q}_0-u_{0,0}\mathcal{Q}_0\otimes\mathcal{L}_0,\\ &d_0(\mathcal{G}_1)=u_{0,0}\mathcal{L}_1\otimes\mathcal{Q}_0-u_{0,0}\mathcal{Q}_0\otimes\mathcal{L}_1,\ d_0(\mathcal{G}_{-1})=u_{0,0}\mathcal{L}_{-1}\otimes\mathcal{Q}_0-u_{0,0}\mathcal{Q}_0\otimes\mathcal{L}_{-1}.\end{aligned}$$

Applying d_0 to $[\mathcal{L}_{-1},\mathcal{Q}_1]=-\mathcal{Q}_0$, we have $\mathcal{L}_{-1}\cdot d_0(\mathcal{Q}_1)+d_0(\mathcal{Q}_0)=(-1)^{[d]}\mathcal{Q}_1\cdot d_0(\mathcal{L}_{-1})$. Comparing the coefficients of $\mathcal{L}_i\otimes\mathcal{H}_{-i}$, we obtain $(i+2)A_{1,i}=(i-1)A_{1,i-1}$, $\forall\ i\neq0,1,2$ and then $A_{1,i}=A_{1,-1}+2A_{1,0}=A_{1,-2}-A_{1,0}=0$, $\forall\ i\neq-1,-2,0$. Similarly, comparing the coefficients of $\mathcal{L}_i\otimes\mathcal{Q}_{-i}$, $\mathcal{G}_i\otimes\mathcal{H}_{-i}$, $\mathcal{G}_i\otimes\mathcal{Q}_{-i}$, we obtain

$$x_{1,i}=x_{1,-1}+2x_{1,0}=x_{1,-2}-x_{1,0}=0\ \text{for}\,x=D,E^{\dagger},R,\forall\ i\neq-1,-2,0.$$

Comparing the coefficients of $\mathcal{H}_i\otimes\mathcal{L}_{-i}$, one has $(i+1)A^{\dagger}_{1,i}=(i-2)A^{\dagger}_{1,i-1}$, $\forall\ i\neq-1,-2,0$, which implies. $A^{\dagger}_{1,i}=2A^{\dagger}_{1,-1}+A^{\dagger}_{1,0}=2A^{\dagger}_{1,1}+A^{\dagger}_{1,0}=0$, $\forall\ i\neq-1,1,0$. Similarly comparing the coefficients of $\mathcal{Q}_i\otimes\mathcal{L}_{-i}$, $\mathcal{H}_i\otimes\mathcal{G}_{-i}$, $\mathcal{Q}_i\otimes\mathcal{G}_{-i}$, we have

$$x_{1,i}=2x_{1,-1}+x_{1,0}=2x_{1,1}+x_{1,0}=0,\ \text{for}\,x=D^{\dagger},E,R^{\dagger},\forall\ i\neq-1,1,0.$$

Comparing the coefficients of $\mathcal{L}_i\otimes\mathcal{G}_{-i}$, one has $(i+2)B_{1,i}=(i-2)B_{1,i-1}$ and

$$B_{1,i}=B_{1,-1}+B_{1,0}=3B_{1,1}+B_{1,0}=3B_{1,-2}-B_{1,0}=0,\ \forall\ i\neq-1,-2,0,1.$$

Comparing the coefficients of $\mathcal{G}_i\otimes\mathcal{L}_{-i}$, $\mathcal{L}_i\otimes\mathcal{L}_{-i}$, $\mathcal{G}_i\otimes\mathcal{G}_{-i}$, one has

$$x_{1,i}=x_{1,-1}+x_{1,0}=3x_{1,1}+x_{1,0}=3x_{1,-2}-x_{1,0}=0,\ \text{for}\,x=X,M,B^{\dagger},\forall\ i\neq-1,-2,0,1.$$

Comparing the coefficients of $\mathcal{H}_i \otimes \mathcal{Q}_{-i}$, we obtain $(i+1)F_{1,i} = (i-1)F_{1,i-1}$, $\forall\ i \neq -1, -2, 0, 1, 2$ and $F_{1,i} = F_{1,-1} + F_{1,0} = 0$, $\forall\ i \neq -1, 0$. Similarly, comparing the coefficients of $\mathcal{Q}_i \otimes \mathcal{H}_{-i}$, $\mathcal{H}_i \otimes \mathcal{H}_{-i}$, $\mathcal{Q}_i \otimes \mathcal{Q}_{-i}$, we have $x_{1,i} = x_{1,-1} + x_{1,0} = 0$ for $x = F^\dagger, Y, N$, $\forall\ i \neq -1, 0$. Applying d_0 to $[\mathcal{H}_0, \mathcal{Q}_1] = -\mathcal{Q}_1$, we have $\mathcal{H}_0 \cdot d_0(\mathcal{Q}_1) + d_0(\mathcal{Q}_1) = (-1)^{[d]}\mathcal{Q}_1 \cdot d_0(\mathcal{H}_0)$. Comparing the coefficients of $\mathcal{G}_1 \otimes \mathcal{H}_0$, $\mathcal{H}_0 \otimes \mathcal{G}_1$, $\mathcal{L}_1 \otimes \mathcal{G}_0$, $\mathcal{G}_1 \otimes \mathcal{L}_0$, $\mathcal{G}_1 \otimes \mathcal{G}_0$, $\mathcal{G}_1 \otimes \mathcal{Q}_0$, $\mathcal{Q}_0 \otimes \mathcal{Q}_1$, $\mathcal{L}_1 \otimes \mathcal{H}_0$, $\mathcal{G}_1 \otimes \mathcal{Q}_0$, $\mathcal{H}_1 \otimes \mathcal{L}_0$, $\mathcal{Q}_1 \otimes \mathcal{G}_0$, one has

$$E^\dagger_{1,0} = E_{1,0} = B_{1,0} = B^\dagger_{1,0} = M_{1,0} = A^\dagger_{1,0} = X_{1,0} = A_{1,0} = R_{1,0} = R^\dagger_{1,0} = Y_{1,0} = N_{1,0} = 0.$$

Applying d_0 to $[\mathcal{G}_{-1}, \mathcal{Q}_1] = 2\mathcal{L}_0 - 2\mathcal{H}_0$, we have $(-1)^{[d]}\mathcal{G}_{-1} \cdot d_0(\mathcal{Q}_1) + (-1)^{[d]}\mathcal{Q}_1 \cdot d_0(\mathcal{G}_{-1}) + 2d_0(\mathcal{H}_0) = 0$. Comparing the coefficients of $\mathcal{G}_0 \otimes \mathcal{Q}_0$, $\mathcal{Q}_1 \otimes \mathcal{G}_{-1}$, one has $2D_{1,0} + F_{1,0} = D^\dagger_{1,0} + F^\dagger_{1,0} = 0$. Applying d_0 to $[\mathcal{L}_1, \mathcal{Q}_{-1}] = \mathcal{Q}_0$, we have

$$\mathcal{L}_1 \cdot d_0(\mathcal{Q}_{-1}) = d_0(\mathcal{Q}_0). \tag{3.8}$$

Comparing the coefficients of $\mathcal{L}_i \otimes \mathcal{H}_{-i}$ in (3.8), we obtain $(i-2)A_{-1,i} = (i+1)A_{-1,i+1}$, $\forall\ i \neq 0, 1$ and $A_{-1,i} = A_{-1,1} + 2A_{-1,0} = A_{-1,2} - A_{-1,0} = 0$, $\forall\ i \neq 1, 0, 2$. Comparing the coefficients of $\mathcal{L}_i \otimes \mathcal{Q}_{-i}$, $\mathcal{G}_i \otimes \mathcal{H}_{-i}$, $\mathcal{G}_i \otimes \mathcal{Q}_{-i}$, we have

$$x_{-1,i} = x_{-1,1} + 2x_{-1,0} = x_{-1,2} - x_{-1,0} = 0 \text{ for } x = D, E^\dagger, R,\ \forall\ i \neq 1, 0, 2.$$

Comparing the coefficients of $\mathcal{H}_i \otimes \mathcal{L}_{-i}$ in (3.8), one has $A^\dagger_{-1,1} = A^\dagger_{-1,-1} = -\frac{1}{2}A^\dagger_{-1,0}$, $\forall\ i \neq 1, -1, 0$. Comparing the coefficients of $\mathcal{Q}_i \otimes \mathcal{L}_{-i}$, $\mathcal{H}_i \otimes \mathcal{G}_{-i}$, $\mathcal{Q}_i \otimes \mathcal{G}_{-i}$, we have

$$x_{-1,i} = 0,\ x_{-1,1} = x_{-1,-1} = -\frac{1}{2}x_{-1,0} \text{ for } x = D^\dagger, E, R^\dagger, \forall\ i \neq 1, -1, 0.$$

Comparing the coefficients of $\mathcal{L}_i \otimes \mathcal{G}_{-i}$, we obtain $(i-2)B_{-1,i} = (i+2)B_{-1,i+1}$ for $i \in \mathbb{Z}$ and

$$B_{-1,i} = B_{-1,1} + B_{-1,0} = 3B_{-1,-1} + B_{-1,0} = 3B_{-1,-2} - B_{-1,0} = 0,\ \forall\ i \neq 1, -1, -2, 0.$$

Comparing the coefficients of $\mathcal{G}_i \otimes \mathcal{L}_{-i}$, $\mathcal{L}_i \otimes \mathcal{L}_{-i}$, $\mathcal{G}_i \otimes \mathcal{G}_{-i}$, one has $x_{-1,i} = x_{-1,1} + x_{-1,0} = 3x_{-1,-1} + x_{-1,0} = 3x_{-1,-2} - x_{-1,0} = 0$ for $x = X, M, B^\dagger$, $\forall\ i \neq 1, -1, -2, 0$. Comparing the coefficients of $\mathcal{H}_i \otimes \mathcal{Q}_{-i}$ in (3.8), we have $(i-1)F_{-1,i} = (i+1)F_{-1,i+1}$, $\forall\ i \neq 0$ and $F_{-1,i} = F_{-1,1} + F_{-1,0} = 0$, $\forall\ i \neq 1, 0$. Comparing the coefficients of $\mathcal{Q}_i \otimes \mathcal{H}_{-i}$, $\mathcal{H}_i \otimes \mathcal{H}_{-i}$, $\mathcal{Q}_i \otimes \mathcal{Q}_{-i}$, one has $x_{-1,i} = x_{-1,1} + x_{-1,0} = 0$ for $x = F^\dagger, Y, N$, $\forall\ i \neq 1, 0$.

Applying d_0 to $[\mathcal{H}_0, \mathcal{Q}_{-1}] = \mathcal{Q}_{-1}$, we have $\mathcal{H}_0 \cdot d_0(\mathcal{Q}_{-1}) = (-1)^{[d]}\mathcal{Q}_{-1} \cdot d_0(\mathcal{H}_0) - d_0(\mathcal{Q}_{-1})$. Comparing the coefficients of $\mathcal{G}_{-1} \otimes \mathcal{H}_0$, $\mathcal{L}_{-1} \otimes \mathcal{G}_0$, $\mathcal{G}_{-1} \otimes \mathcal{L}_0$, $\mathcal{G}_{-1} \otimes \mathcal{G}_0$, $\mathcal{Q}_0 \otimes \mathcal{Q}_{-1}$, $\mathcal{L}_{-1} \otimes \mathcal{H}_0$, $\mathcal{H}_{-1} \otimes \mathcal{H}_0$, $\mathcal{G}_{-1} \otimes \mathcal{Q}_0$, $\mathcal{Q}_{-1} \otimes \mathcal{G}_0$, $\mathcal{H}_{-1} \otimes \mathcal{L}_0$, $\mathcal{L}_{-1} \otimes \mathcal{L}_0$, one has $E_{-1,0} = E^\dagger_{-1,0} = B_{-1,0} = B^\dagger_{-1,0} = M_{-1,0} = R_{-1,0} = R^\dagger_{-1,0} = A^\dagger_{-1,0} = X_{-1,0} = N_{-1,0} = A_{-1,0} = Y_{-1,0} = 0$. Applying

d_0 to $[\mathcal{L}_2, \mathcal{Q}_{-1}] = \mathcal{Q}_1$, we have $\mathcal{L}_2 \cdot d_0(\mathcal{Q}_{-1}) = (-1)^{[d]}\mathcal{Q}_{-1} \cdot d_0(\mathcal{L}_2) + d_0(\mathcal{Q}_1)$. Comparing the coefficients of $\mathcal{L}_2 \otimes \mathcal{Q}_{-1}$, $\mathcal{L}_1 \otimes \mathcal{Q}_2$, $\mathcal{Q}_{-2} \otimes \mathcal{L}_3$, $\mathcal{Q}_2 \otimes \mathcal{L}_{-1}$, $\mathcal{H}_1 \otimes \mathcal{Q}_0$, $\mathcal{Q}_1 \otimes \mathcal{H}_0$, one has

$$D_{-1,0} = D_{1,0} = D^{\dagger}_{-1,0} = D^{\dagger}_{1,0} = F_{-1,0} - F_{1,0} = F^{\dagger}_{-1,0} - F^{\dagger}_{1,0} = 0.$$

Then $F^{\dagger}_{1,0} = F_{1,0} = F^{\dagger}_{-1,0} = F_{-1,0} = 0$ and $d_0(\mathcal{Q}_1) = d_0(\mathcal{Q}_{-1}) = 0$.

Applying d_0 to $[\mathcal{L}_1, \mathcal{H}_{-1}] = \mathcal{H}_0$, we have $\mathcal{L}_1 \cdot d_0(\mathcal{H}_{-1}) = d_0(\mathcal{H}_0)$. Comparing the coefficients of $\mathcal{L}_i \otimes \mathcal{H}_{-i}$, we have $(i-2)\xi_{-1,i} = (i+1)\xi_{-1,i+1}$ for $i \in \mathbb{Z}$. we obtain $\xi_{-1,i} = \xi_{-1,1} + 2\xi_{-1,0} = \xi_{-1,2} - \xi_{-1,0} = 0$, $\forall\ i \neq 0, 1, 2$. Similarly comparing the coefficients of $\mathcal{L}_i \otimes \mathcal{Q}_{-i}$, $\mathcal{G}_i \otimes \mathcal{Q}_{-i}$, $\mathcal{L}_i \otimes \mathcal{Q}_{-i}$, we have $x_{-1,i} = x_{-1,1} + 2x_{-1,0} = x_{-1,2} - x_{-1,0} = 0$ for $x = \delta^{\dagger}, \varphi, \varepsilon, \forall\ i \neq 0, 1, 2$. Comparing the coefficients of $\mathcal{H}_i \otimes \mathcal{L}_{-i}$, we have $(i-1)\xi^{\dagger}_{-1,i} = (i+2)\xi^{\dagger}_{-1,i+1}$ for $i \in \mathbb{Z}$ and

$$\xi^{\dagger}_{-1,i} = 0,\ \xi^{\dagger}_{-1,1} = \xi^{\dagger}_{-1,-1} = -\frac{1}{2}\xi^{\dagger}_{-1,0},\ \forall\ i \neq 1, -1, 0.$$

Comparing the coefficients of $\mathcal{H}_i \otimes \mathcal{G}_{-i}$, $\mathcal{Q}_i \otimes \mathcal{G}_{-i}$, $\mathcal{Q}_i \otimes \mathcal{L}_{-i}$, we have $x_{-1,i} = 0$, $x_{-1,1} = x_{-1,-1} = -\frac{1}{2}x_{-1,0}$ for $x = \delta, \varphi^{\dagger}, \varepsilon^{\dagger}, \forall\ i \neq 1, -1, 0$. Comparing the coefficients of $\mathcal{L}_i \otimes \mathcal{G}_{-i}$, we have $(i-2)\eta_{-1,i} = (i+2)\eta_{-1,i+1}$ for $i \in \mathbb{Z}$ and

$$\eta_{-1,i} = \eta_{-1,1} + \eta_{-1,0} = 3\eta_{-1,-1} + \eta_{-1,0} = 3\eta_{-1,2} - \eta_{-1,0} = 0,\ \forall\ i \neq 1, -1, 2, 0.$$

Comparing the coefficients of $\mathcal{G}_i \otimes \mathcal{L}_{-i}$, $\mathcal{L}_i \otimes \mathcal{L}_{-i}$, $\mathcal{G}_i \otimes \mathcal{G}_{-i}$, we have

$$x_{-1,i} = x_{-1,1} + x_{-1,0} = 3x_{-1,-1} + x_{-1,0} = 3x_{-1,2} - x_{-1,0} = 0,\ x = \eta^{\dagger}, h, g,\ \forall\ i \neq 0, \pm 1, 2.$$

Comparing the coefficients of $\mathcal{H}_i \otimes \mathcal{Q}_{-i}$, we obtain $(i-1)\psi_{-1,i} = (i+1)\psi_{-1,i+1}$ for $i \in \mathbb{Z}^*$ and $\psi_{-1,i} = \psi_{-1,1} + \psi_{-1,0} = 0$, $\forall\ i \neq 1, 0$. Comparing the coefficients of $\mathcal{Q}_i \otimes \mathcal{H}_{-i}$, $\mathcal{H}_i \otimes \mathcal{H}_{-i}$, we have $x_{-1,i} = x_{-1,1} + x_{-1,0} = 0$ for $x = \psi^{\dagger}, l, \forall\ i \neq 1, 0$. According to $\mathcal{Q}_i \otimes \mathcal{Q}_{-i}$, we obtain $(i-1)v_{-1,i} = (i+1)v_{-1,i+1}$ for $i \neq 0$ and $v_{-1,i} = v_{-1,1} + v_{-1,0} - (-1)^{[d]+1}u_{0,0} = 0$, $\forall\ i \neq 1, 0$.

Applying d_0 to $[\mathcal{H}_{-1}, \mathcal{G}_1] = \mathcal{G}_0$, we have $\mathcal{H}_{-1} \cdot d_0(\mathcal{G}_1) = (-1)^{[d]}\mathcal{G}_1 \cdot d_0(\mathcal{H}_{-1}) + d_0(\mathcal{G}_0)$. Comparing the coefficients of $\mathcal{G}_1 \otimes \mathcal{Q}_{-1}$, $\mathcal{L}_0 \otimes \mathcal{L}_0$, one has $2\varepsilon_{-1,0} - \psi_{-1,0} = 2\varepsilon_{-1,0} - \varepsilon^{\dagger}_{-1,0} = 0$. Applying d_0 to $[\mathcal{H}_{-1}, \mathcal{Q}_1] = -\mathcal{Q}_0$, we have $\mathcal{H}_{-1} \cdot d_0(\mathcal{Q}_1) + d_0(\mathcal{Q}_0) = (-1)^{[d]}\mathcal{Q}_1 \cdot d_0(\mathcal{H}_{-1})$. Comparing the coefficients of $\mathcal{H}_1 \otimes \mathcal{Q}_{-1}$, $\mathcal{Q}_1 \otimes \mathcal{H}_{-1}$, $\mathcal{Q}_0 \otimes \mathcal{Q}_0$, $\mathcal{Q}_2 \otimes \mathcal{Q}_{-2}$, $\mathcal{Q}_2 \otimes \mathcal{H}_{-2}$, $\mathcal{H}_{-2} \otimes \mathcal{Q}_2$, $\mathcal{H}_1 \otimes \mathcal{H}_{-1}$, $\mathcal{H}_{-1} \otimes \mathcal{H}_1$, $\mathcal{Q}_{-2} \otimes \mathcal{L}_2$, $\mathcal{L}_{-2} \otimes \mathcal{L}_2$, $\mathcal{L}_2 \otimes \mathcal{L}_{-2}$, $\mathcal{Q}_{-1} \otimes \mathcal{Q}_1$, we have $\varepsilon_{-1,0} = l_{-1,0} = \xi_{-1,0} = \xi^{\dagger}_{-1,0} = \delta_{-1,0} = \delta^{\dagger}_{-1,0} = \eta_{-1,0} = \eta^{\dagger}_{-1,0} = \varphi^{\dagger}_{-1,0} = h_{-1,0} = g_{-1,0} = \varphi_{-1,0} = \varepsilon^{\dagger}_{-1,0} = \psi_{-1,0} = \psi^{\dagger}_{-1,0} = 0$. Applying d_0 to $[\mathcal{H}_{-1}, \mathcal{H}_0] = 0$, we have $\mathcal{H}_{-1} \cdot d_0(\mathcal{H}_0) = \mathcal{H}_0 \cdot d_0(\mathcal{H}_{-1})$. Comparing the coefficients of $\mathcal{Q}_0 \otimes \mathcal{Q}_{-1}$, one has $u_{0,0} = 0$. Then $d_0(\mathcal{G}_{\pm 1}) = d_0(\mathcal{G}_0) = d_0(\mathcal{H}_0) = d_0(\mathcal{H}_{-1}) = 0$.

Claim 3. The sum in (3.1) is finite.

By now, we have completed the proof of Proposition 3.2.

The following lemma can be obtained by using the similar techniques developed in [5].

Lemma 3.3. *If* $x \cdot r \in \mathrm{Im}(1 \otimes 1 - \tau)$ *for some* $r \in V$ *and any* $x \in \mathcal{T}$, *then* $r \in \mathrm{Im}(1 \otimes 1 - \tau)$.

Proof of Theorem 1.1. Suppose $(\mathcal{T}, [\cdot,\cdot], \triangle)$ is a Lie super-bialgebra structure on $\mathcal{T}$. Then $\Delta = \Delta_r$ for some $r \in (\mathcal{T} \otimes \mathcal{T})^{\bar{0}}$ by Proposition 3.2. Then $\mathrm{Im}\,\Delta \subset \mathrm{Im}(1 \otimes 1 - \tau)$. Thus by Lemma 3.3, we have $r \in \mathrm{Im}(1 \otimes 1 - \tau)$. Then Lemma 3.1(iii) shows that $c(r) = 0$. Hence $(\mathcal{T}, [\cdot,\cdot], \triangle)$ is a triangular coboundary Lie super-bialgebra.

References

[1] G. R. Beatriz, Construction formulae for singular vectors of the topological and of the Ramond $N=2$ superconformal algebras, *Int. J. Mod. Phys.*, **17** (2002), 4515–4541.

[2] V.G. Kac, Lie superalgebras, *Adv. Math.*, **26** (1977), 8–97.

[3] S.H. Ng and E.J. Taft, Classification of the Lie bialgebra structures on the Witt and Virasoro algebras, *J. Pure Appl. Alg.*, **151** (2000), 67–88.

[4] E.J. Taft, Witt and Virasoro algebras as Lie bialgebras, *J. Pure Alg.*, **87** (1993), 301–312.

[5] H. Yang and Y. Su, Lie super-bialgebra structures on the Ramond $N = 2$ super-virasoro algebra, *Chaos, Solitons and Fractals*, 2008.

[6] H. Fa and J. Li, Lie super-bialgebra structures on the centerless twisted $N=2$ superconformal algebra, submitted.